

RESEARCH ARTICLE

ANTIOXIDANT ACTIVITY OF ETHANOLIC EXTRACT OF *CASSIA HIRSUTA* (L.) LEAVES

Tapas Kanti Dey^{1*}, Talha Bin Emran¹, Dibyajyoti Saha¹, Md. Atiar Rahman², S.M. Zahid Hosen¹ and Nipa Chowdhury¹

¹Department of Pharmacy, BGC Trust University, Chittagong, Bangladesh

²Department of Biochemistry and Molecular Biology, University of Chittagong, Chittagong, Bangladesh

*E-mails: tapas85@yahoo.com, saha.dibyajyoti@gmail.com

Tel.: +88-01815520598, +88-01819942214.

Received: March 21, 2012 / Revised: June 15, 2012 / Accepted: June 16, 2012

The antioxidant activity of ethanolic extract of *Cassia hirsuta* (Family-Caesalpinaceae) leaves has been investigated in the present study. The antioxidant activity of the *Cassia hirsuta* leaves ethanolic extract was assessed by DPPH (1, 1-diphenyl-2-picrylhydrazyl) free radical scavenging method. Extract showed significant DPPH free radical scavenging effect compared with standard antioxidant ascorbic acid. IC₅₀ value of ascorbic acid and leaves extract was found 1.25 µg/ml and 200.96 µg/ml, respectively. The value of extract indicated significant antioxidant activity of the plant.

Key words: *Cassia hirsuta*, Antioxidant, DPPH, *Caesalpinaceae*.

INTRODUCTION

Antioxidants are type of molecules that neutralize harmful free radicals, produced through a chain of reactions that damage living cells, spoil foods; degrade materials such as rubber, gasoline, lubricating oil. Antioxidants terminate these chain reactions through the removal of free radical intermediates and inhibition of other oxidation reactions (Sies, 1997). This is why plants and animals maintain complex systems of multiple antioxidants, such as glutathione, vitamin C, and vitamin E along with some enzymes like catalase, superoxide dismutase and various peroxidases. The use of antioxidants in pharmacology is intensively studied as oxidative stress might be an important part of many human diseases particularly stroke and neurodegenerative incidents (Joseph *et al* 2010). Antioxidants, therefore, are routinely added to meals, oils, foodstuffs, and other materials to prevent free radical damage. Recently there has been an upsurge of interest in the therapeutic potentials of medicinal plants as antioxidants in reducing such free radical induced tissue injury. A lot of new plant species have been investigated in the

search for novel antioxidants (Chu *et al* 2000; Koleva *et al* 2002; Mantle *et al* 2000; Oke *et al* 2002) other than well known and traditionally used natural antioxidants from tea, wine, fruits, vegetables and spices (Schuler, 1990) but there is still a demand to find more information on the antioxidant potential of plant species.

Cassia hirsuta Linn. plant (Irwin and Barneby, 1982; Holm *et al* 1979) belongs to the family *Caesalpinaceae* and commonly called as stinking cassia and hairy senna (**Figure 1**).



Fig. 1. Photograph of *Cassia hirsuta* Linn.

It is a terrestrial perennial, erect shrub up to 150 cm in tall, stem rounded, solid glabrous, flowering period from September to December and fruiting in November to January. It is a native of tropical America and now distributed in Malaysia, Indo-China, Thailand, Asian and African tropics, Laos, Java, Brazil, California, New Mexico and India (Holm *et al* 1979). It is used as a green manure and forage plant. In Africa, it planted as a shade plant in young coffee plantation. Leaves and young pods are eaten, usually steamed or cooked in vegetable or in salads. In Java, the leaves are used medicinally for treating herpes. A decoction of leaves is used against irritation of skin in Thailand. In Laos, the seeds are used as a substitute for coffee. Phytomedicinally, the plant parts or extracts are used for treating illness in man. Plants as gifts of nature have many therapeutic properties combined with much nutritive value, which have made their use in chemotherapy as valuable as the synthetic drugs. Herbal organ of the body are used to feed and restore to health those parts, which have become weakened. It is a medicinal plant widely used for stomach troubles, dysentery, abscesses, rheumatism, fever and other diseases. Seeds contain phytotoxin, tannins and 0.25% chrysoarobin. Seeds also contain a water soluble sugars extract as D-galactose and D-mannose in 1:4 molar ratio from hydrolysed compound on paper chromatogram Present manuscript mainly deals with the methylation studies for polysaccharide structure of *Cassia hirsuta* Linn. seeds - galactomannan. It is also useful in dental caries: powdered seeds are used to massage on gums and teeth (Brenan, 1967; Revathi and Parimelazhagan, 2010; Jamir *et al* 2010; Singh *et al* 2007).

Scientists have been proving that all the natural things are not good for health. Different retrospective studies done over the last 20 years indicated that the incidence of deaths occurring due to exposure to plants (as a proportion of total patients poisoned by traditional plant medicine) was about 1.5% in France, 5% in Belgium, 6.5% in Italy, 7.2% in Switzerland and 6% in Turkey (Gaillard *et al* 1999). Therefore, there is a need to have an understanding of the risks posed by herbal medicines so as to ensure that such products could be used safely.

Keeping in view the biopotential of plants and their derived products (Dahiya and Gautam, 2011; Jain *et al* 2011; Zahid Hosen *et al* 2011; Chowdhury *et al* 2012; Emran *et al* 2012), present investigation was directed toward

evaluating the antioxidant property of *Cassia hirsuta* leaves extract.

MATERIALS AND METHODS

Collection of plant

The plant *Cassia hirsuta* was collected from the Chittagong University campus. The plant was taxonomically classified and identified scientifically by Professor Dr. Md. Mostafa Kamal Pasha, Taxonomist, Department of Botany, University of Chittagong, Bangladesh with help of direct field study and taxonomic analysis.

Preparation of plant extract

The fresh leaves of *Cassia hirsuta* were washed with water immediately after collection. Then, these were chopped into small pieces, air dried at room temperature for about 10 days and pulverized into powder form and stored in an airtight container. 600 g leaf powder was macerated in 6 litre pure ethanol for 7 days at room temperature. After 7 days, leaf ethanol extract was filtered off through a cotton plug and finally with a Whatman No. 1 filter paper. Then the extract was concentrated under reduced pressure below 50 °C through rotary vacuum evaporator. The concentrated extracts were collected in a petri dish and allowed to air dry for complete evaporation of ethanol. The whole process were repeated three times and finally, 72 g blackish-green colored, concentrated leaf extract was obtained (yield 12% w/w) which was kept in refrigerator at 4 °C.

In vitro assay for antioxidant activity of plant extract

The antioxidant activity of *Cassia hirsuta* leaves extract was assessed in comparison to standard antioxidant ascorbic acid (BDH, England) depending on the scavenging effect of 1, 1-diphenyl-2-picrylhydrazyl (DPPH)-free radical. The whole procedure was done according to established procedure (Braca *et al* 2001; Brand-William *et al* 1995). Ascorbic acid solution (5 ml) and different concentrations of extract (20, 40, 60, 80, 100, 200, 400 and 800 µg/ml in methanol) (5 ml) were mixed with 3 ml of 0.4 mM (0.004 %) DPPH solution. The mixtures were kept in dark for 30 min to measure the absorbance at 517 nm using UV-Visible spectrophotometer (UV-1601 Shimadzu, Kyoto, Japan) and ascorbic acid was used as a positive control. Lower absorbance of the reaction mixture indicated higher free radical-scavenging activity. The degree of decolorization of DPPH

from purple to yellow indicated the scavenging efficiency of the extract. The scavenging activity against DPPH was calculated using the following equation:

$$\text{Scavenging activity (\%)} = [(A - B) / A] \times 100 \quad (1)$$

Where A is absorbance of control (DPPH solution without the sample) and B is the absorbance of DPPH solution in the presence of the sample (extract/ascorbic acid). The scavenging activity (%) or % inhibition was then plotted against log

concentration and from the graph IC₅₀ (Inhibition concentration 50) value was calculated by linear regression analysis.

RESULTS AND DISCUSSION

Assay for antioxidative activity of Cassia hirsuta

DPPH free radical scavenging activity of *Cassia hirsuta* and ascorbic acid is shown in the **Table 1**. Both ascorbic acid and *Cassia hirsuta* leaves ethanolic extract showed dose dependent activity.

Table 1. DPPH free radical scavenging activity of ascorbic acid and *Cassia hirsuta* leaves extract

Test material	Concentration ($\mu\text{g/ml}$)	% Scavenging activity	IC ₅₀ ($\mu\text{g/ml}$)
Ascorbic acid (Standard)	20	70.02	IC ₅₀ = 1.25
	40	74.06	
	60	79.25	
	80	84.09	
	100	88.50	
	200	90.75	
	400	94.50	
	800	97.34	
<i>Cassia hirsuta</i>	20	7.04	IC ₅₀ = 200.96
	40	17.25	
	60	25.70	
	80	27.11	
	100	31.51	
	200	54.40	
	400	63.20	
	800	78.70	

Among the eight different concentrations used in the study (20, 40, 60, 80, 100, 200, 400 and 800 $\mu\text{g/ml}$) ascorbic acid showed 70.02, 74.06, 79.25, 84.09, 88.50, 90.75, 94.50 and 97.34% scavenging activity where highest scavenging activity was 97.34% at concentration 800 $\mu\text{g/ml}$. On the other hand, *Cassia hirsuta* leaf ethanol extract showed 7.04, 17.25, 25.70, 27.11, 31.51, 54.40, 63.20 and 78.70% scavenging activity at the above mentioned eight different concentrations where highest scavenging activity of *Cassia hirsuta* leaf ethanol extract was 78.70% at 800 $\mu\text{g/ml}$.

% of scavenging activity or % of inhibition was plotted against log concentration and from the graph IC₅₀ (Inhibition concentration 50) value was calculated by linear regression analysis.

IC₅₀ value of ascorbic acid and *Cassia hirsuta* leaves ethanolic extract was found 1.25 and 200.96 $\mu\text{g/ml}$ respectively (**Figure 2**). The IC₅₀

value obtained for *Cassia hirsuta* extract and ascorbic acid indicated that *Cassia hirsuta* extract possessed higher efficiency to neutralize free radicals than that of ascorbic acid (**Figure 3**). Different research suggested that most of the plant extracts showing antioxidant activity are due to presence of the phenolic compounds (Ramarathnam *et al* 1997; Velioglu *et al* 1998). Phenolic natural compounds such as flavonoids possess antioxidant activity due to their redox properties which allow them to act as reducing agents and singlet oxygen quencher. In addition, they have metal chelating potentials (Rice-Evans *et al* 1995). The phenolic compounds, identified in the extract might contribute to the antioxidant activity of *Cassia hirsuta* extract.

CONCLUSION

The results of the study demonstrated that the ethanolic extract of *Cassia hirsuta* exhibited very

potential antioxidant effect in experimental models which supported the claims by traditional medicine practitioners. These results can be strong scientific evidence to use this plant

as a useful source of antioxidant references. However, further studies are still necessary to elucidate a mechanistic way how the plant contributes in these pharmacologic properties.




Fig. 2. Comparative IC₅₀ values of reference antioxidant and studied plant *Cassia hirsuta* leaf extract: (a) ascorbic acid (b) plant extract




Fig. 3. Comparative % scavenging activities of *Cassia hirsuta* extract and ascorbic acid

REFERENCES

- Braca A, De Tommasi N, Di Bari L, Pizza C, Politi M, Morelli I. Antioxidant principles from *Bauhinia tarapotensis*. *J. Nat. Prod.* 2001;64(7):892-5. [DOI: 10.1021/np0100845]
- Brand-William W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT - Food Sci. Technol.* 1995;28(1):25-30. [DOI: 10.1016/S0023-6438(95)80008-5]
- Brenan JPM. Leguminosae, subfamily Caesalpinioideae. In: Milne-Redhead E, Polhill RM. (eds). *Flora of Tropical East Africa*. Crown Agents for overseas governments and administrations: London, UK, 1967; 230.
- Chowdhury S, Saha D, Paul S. *In vitro* cytotoxic activities of methanolic extract of *Mimosa pudica*. *Bull. Pharm. Res.* 2012;2(1):42-5.
- Chu Y-H, Chang C-L, Hsu-H-F. Flavonoid content of several vegetables and their antioxidant activity. *J. Sci. Food Agr.* 2000;80(5):561-6.
- Dahiya R, Gautam H. Solution phase synthesis and bioevaluation of cordyheptapeptide B. *Bull. Pharm. Res.* 2011;1(1):1-10.
- Emran TB, Rahman MA, Zahid Hosen SM, Saha D, Chowdhury S, Saha D, Dey TK. Antioxidant property of ethanolic extract of *Leucas aspera* Linn. *Bull. Pharm. Res.* 2012;2(1):46-9.
- Holm LG, Pancho JV, Herberger JP, Plucknett DL. *A Geographical Atlas of World Weeds*, Knieger Publishing Company: Florida, 1979.
- Irwin HS, Barneby RC. The American Cassiinae, A Synoptical revision of Leguminosae tribe Cassieae subtribe Cassiinae in the New World. *Mem. New York Bot. Gar.* 1982;35:435.
- Jain RA, Agarwal RC, Pandey A, Jain R. Evaluation of *Argemone mexicana* fruits extract using micronucleus assay in mouse bone marrow cells. *Bull. Pharm. Res.* 2011;1(2):22-4.

- Jamir NS, Takatemjen, Limasemba. Traditional knowledge of *Lotha-Naga* tribes in Wokha district, Nagaland. *Ind. J. Trad. Knowl.* 2010;9(1):45-8.
- Jenny A, Saha D, Paul S, Dutta M, Uddin MZ, Nath AK. Antibacterial activity of aerial part of extract of *Elephantopus scaber* Linn. *Bull. Pharm. Res.* 2012;2(1):38-41.
- Joseph NM, Sabharwal M, Shashi A, Mahor A, Rawal S. *In-vitro* and *in-vivo* models for antioxidant activity evaluation - a review. *Int. J. Pharm. Sci. Res.* 2010;1(1):1-11.
- Koleva II, van Beek TA, Linssen JPH, de Groot A, Evstatieva LN. Screening of plant extracts for antioxidant activity: a comparative study on three testing methods. *Phytochem. Anal.* 2002;13(1):8-17. [DOI: 10.1002/pca.611]
- Mantle D, Eddeb F, Pickering AT. Comparison of relative antioxidant activities of British medicinal plant species in vitro. *J. Ethnopharmacol.* 2000;72(1-2):47-51. [DOI: 10.1016/S0378-8741(00)00199-9]
- Oke JM, Hamburger MO. Screening of some nigerian medicinal plants for antioxidant activity using 2, 2, diphenyl-picryl-hydrazyl radical. *Afr. J. Biomed. Res.* 2002; 5(1-2):77-9.
- Ramarathnam N, Ochi H, Takeuchi M. Antioxidant Defense System in Vegetable Extracts. In: Shahidi F. (ed.) *Natural Antioxidants: Chemistry, Health Effects and Applications*, Aocs Press: Champaign, IL, USA, 1997; 76-87.
- Revathi P, Parimelazhagan T. Traditional knowledge on medicinal plants used by the Irula Tribe of Hasanur Hills, Erode district, Tamil Nadu, India. *Ethnobot. Leaf.* 2010; 14:136-60.
- Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radic. Res.* 1995;22(4):375-83.
- Schuler P. Natural antioxidants exploited commercially. In: Hudson BJF. (ed.) *Food Antioxidants*. Elsevier: London, 1990; 99-170.
- Sies H. Oxidative stress: oxidants and antioxidants. *Exp. Physiol.* 1997;82(2):291-5.
- Singh J, Kumar A, Budhiraja S, Hooda A. Ethnomedicine: use in dental caries. *Braz. J. Oral Sci.* 2007;6(21):1308-12.
- Velioglu YS, Mazza G, Gao L, Oomah BD. Antioxidant activity and total phenolics in selected fruits and vegetables, and grain products. *J. Agric. Food Chem.* 1998;46(10):4113-7. [DOI: 10.1021/jf9801973]
- Zahid Hosen SM, Das R, Rahim ZB, Chowdhury N, Paul L, Saha D. Study of Analgesic activity of the methanolic extract of *Acorus calamus* L. and *Oroxylum indicum* Vent by acetic acid induced writhing method. *Bull. Pharm. Res.* 2011;1(3):63-7.
<http://www.africamuseum.be/collections>
