

Basarkar GD, Shirsath GN, Patil SB. Development of microspheres containing diclofenac diethylamine as sustained release topical formulation. *Bull. Pharm. Res.* 2013;3(1):14-22.

References (13):

1. Dahiya S, Gupta ON. Formulation and *in vitro* evaluation of metoprolol tartrate microspheres. *Bull. Pharm. Res.* 2011;1(1):31-9.
<http://www.appconnect.in/wp-content/uploads/2011/05/FirstPagePreviewBPR006.pdf>
2. Del Rosso JQ. Benzoyl peroxide microsphere formulation: What is the Science Supporting Microsphere Vehicle Technology and Clinical Use?. *J. Clin. Aesthet. Dermatol.* 2009;2(9):46-54.
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2923969/>
3. Gowda DV, Manjunatha M, Balmurlidhara V, Khan MS. Study on encapsulation of Ranolazine in bees wax microspheres: Preparation, characterization and release kinetics of microspheres. *Der Pharm. Lett.* 2010;2(6):232-43.
<http://scholarsresearchlibrary.com/DPL-vol2-iss6/DPL-2010-2-6-232-243.pdf>
4. Gowda DV, Shivakumar HG. Preparation and evaluation of waxes/fat microspheres loaded with lithium carbonate for controlled release. *Indian J. Pharm. Sci.* 2007;69(2):251-6.
http://www.ijpsonline.com/temp/IndianJPharmSci692251-2547229_070432.pdf
5. Grace D, Rogers J, Skeith K, Anderson K. Topical diclofenac versus placebo: A double blind, randomized clinical trial in patients with osteoarthritis of the knee. *J. Rheumatol.* 1999;26(12):2659-63.
<http://www.ncbi.nlm.nih.gov/pubmed/10606379>
6. Iannuccelli V, Sala N, Tursilli R, Coppi G, Scalia S. Influence of liposphere preparation on butyl-methoxydibenzoylmethane photostability. *Eur. J. Pharm. Biopharm.* 2006;63(2):140-5.
<http://www.ncbi.nlm.nih.gov/pubmed/16531026>
7. Indian Pharmacopoeia, Government of India and Ministry of Family Health Welfare, Vol. I, Indian Pharmacopoeia Commission: Ghaziabad, India, 2007; 80-2.
http://indianpharmacopoeia.in/BookDetail.php?item_id=225&
8. Nasr M, Mansour S, Mortada N, El Shamy AA. Lipospheres as carriers for topical delivery of aceclofenac: Preparation, characterization and *in vivo* evaluation. *AAPS PharmSciTech.* 2008;9(1):154-62. [DOI: 10.1208/s12249-007-9028-2]
<http://www.ncbi.nlm.nih.gov/pubmed/18446476>

9. Nishihata T, Kotera K, Nakano Y, Yamazaki M. Rat percutaneous transport of diclofenac and influence of hydrogenated soya phospholipids. *Chem. Pharm. Bull. (Tokyo)* 1987;35(9):3807-12.
<http://www.ncbi.nlm.nih.gov/pubmed/3435976>
10. Parsaee S, Sarbolouki MN, Parnianpour M. *In-vitro* release of diclofenac diethylammonium from lipid-based formulations. *Int. J. Pharm.* 2002;241(1):185-90.
<http://www.ncbi.nlm.nih.gov/pubmed/12086734>
11. Vali SR, Ju Y-H, Balagopala Kaimal TN, Chern Y-T. A process for the preparation of food-grade rice bran wax and the determination of its composition. *J. Am. Oil Chemists' Soc.* 2005;82(1):57-64.
<http://link.springer.com/article/10.1007%2Fs11746-005-1043-z>
12. Yellanki SK, Singh J, Syed JA, Bigala R, Goranti S, Nerella NK. Design and characterization of amoxicillin trihydrate mucoadhesive microspheres for prolonged gastric retention. *Int. J. Pharm. Sci. Drug Res.* 2010;2(2):112-4.
<http://www.ijpsdr.com/pdf/vol2-issue2/5.pdf>
13. Yokomiza Y. Effect of phospholipids on the percutaneous penetration of drugs through the dorsal skin of the guinea pig, *in vitro*. 3. The effects of phospholipids on several drugs having different polarities. *J. Control. Release* 1996;42(3):217-28.
<http://www.sciencedirect.com/science/article/pii/0168365996013478>

