



RESEARCH ARTICLE

STABILITY-INDICATING UV-VIS SPECTROPHOTOMETRIC METHOD FOR ESTIMATION OF ATORVASTATIN CALCIUM AND FENOFIBRATE IN TABLET DOSAGE FORM

Singh S*, Inamullah, Rai J, Choudhary N, Sharma S

Dept. of Pharmaceutical Chemistry, Invertis Institute of Pharmacy, Invertis University, Bareilly, Uttar Pradesh, India

*E-mail: rssunil29@rediffmail.com

Tel.: +91 8859124788.

Received: November 28, 2012 / Revised: December 23, 2012 / Accepted: December 24, 2012

The present research works discuss the development of a stability-indicating UV spectrophotometric method for the estimation of Atorvastatin calcium (ATC) and Fenofibrate (FEN) in tablet dosages form. The optimum conditions for the analysis of the drug were established. The maximum wavelength (λ_{max}) was found to be 247 nm for ATC and 287 nm for FEN. The linearity of the proposed method was investigated in the range of 6-16 $\mu\text{g/ml}$ and 2-12 $\mu\text{g/ml}$ for ATC, FEN respectively. Calibration curves showed a linear relationship between the absorbance and concentration. The line equation for ATC ($y = 0.041x + 0.043$) with r^2 of 0.999 and for Fenofibrate ($y = 0.054x - 0.003$) with r^2 of 0.999, was obtained. Validation was performed as per ICH guidelines for linearity, accuracy, precision, LOD and LOQ. The LOD was 0.2695 $\mu\text{g/ml}$, 0.0222 $\mu\text{g/ml}$ for ATC and FEN and the LOQ was 0.8780 $\mu\text{g/ml}$, 0.222 $\mu\text{g/ml}$ for ATC and FEN respectively. The proposed method may be suitable for the analysis of ATC and FEN in tablet formulation for quality control purpose. The proposed methods were simple, sensitive, precise, accurate, quick and useful for routine quality control. The stability studies of ATC and FEN were conducted and the degradation characteristics were found to be much more prominent in acid hydrolysis in FEN and alkaline hydrolysis in ATC.

Key words: Simultaneous equation, Degradation, Validation, Atorvastatin calcium, Fenofibrate.

INTRODUCTION

Atorvastatin Calcium (ATC) (Figure 1) is calcium salt of (βR , 8R)-2-(4-fluorophenyl)- α,δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid trihydrate (IP 2007). ATC is a member of the drug class known as statins, used for lowering blood cholesterol. Atorvastatin works by inhibiting HMG-CoA reductase, an enzyme found in liver tissue that plays a key role in production of cholesterol in the body.

Fenofibrate (FEN) (Figure 1) is 2-[4-(4-chloro benzoyl)phenoxy]-2-methyl-propanoic acid, 1-methylethyl ester (ChemSpider, 3222) which is a fibric acid derivative that lowers lipid levels by activating PPAR α and thus, decreases the risk of heart diseases and prevent strokes.




Fig. 1. Structure of ATC and FEN

Atherosclerotic vascular disease is a condition in which an artery wall thickens as a result of accumulation of fatty materials such as cholesterol. It affects mostly arterial blood vessels. Inflammatory response in walls of arteries is commonly referred to as hardening of

arteries. It is caused by formation of multiple plaques with in arteries. Some of drug combination like Atorvastatin calcium and Fenofibrate has a highly beneficial effect on all lipid parameters. Atorvastatin Calcium is more effective in reduction of cholesterol level whereas Fenofibrate is effective in reduction of triglycerides (Diwan *et al* 2012).

Safety and efficacy of pharmaceuticals are two fundamental issues of importance in drug therapy. Instability of pharmaceuticals can cause a change in physical, chemical, pharmacological and toxicological properties of the active pharmaceutical ingredients (API), thereby affecting its safety and efficacy. Hence, the pharmacists should take cognizance of various factors such as drug stability, possible degradation products, mechanisms and routes of degradation and potential interactions with excipients utilized in the formulation to ensure the delivery of their therapeutic values to patients. In order to assess the stability of a drug product, one needs an appropriate analytical methodology, so called the stability indicating methods which allow accurate and precise quantitation of the drug, its degradation products and interaction products, if any (Janardhanan *et al* 2011). Forced degradation studies were performed on Atorvastatin calcium and Fenofibrate to prove the stability indicating property of the method. The stress conditions employed for degradation study included light exposure, acid hydrolysis (0.1 N HCl), base hydrolysis (0.1 N NaOH), and thermal degradation. The duration of time selected for degradation studies was 6 h. The photolytic degradation was performed by exposing the solid drugs to sunlight for 12 h. The concentration of 100 $\mu\text{g/ml}$ of each of Atorvastatin calcium and Fenofibrate were prepared using respective solvents (NaOH, HCl, methanol) separately (ICH Q1A (R2), 2003).

Literature survey revealed that various analytical methods have been reported for estimation of Atorvastatin calcium (ATV) and Fenofibrate (FEB) individually from its formulations and biological fluids (Erturk *et al* 2003; Erk, 2003; Altuntas and Erk, 2004; Gupta *et al* 2010; Salama *et al* 2011). However, no stability-indicating UV spectrophotometric method is reported for the estimation of Atorvastatin calcium (ATC) and Fenofibrate (FEN) in tablet dosages form in the literature.

In continuation of our work on analytical method development of drugs (Singh *et al* 2010;

2011; 2012), the present investigation was directed toward development of a stability-indicating UV spectrophotometric method for the estimation of Atorvastatin calcium and Fenofibrate in tablet dosages form. The objective of the present work is to develop simple, rapid, accurate, specific and economic UV stability indicating method for the estimation of ATC and FEM in bulk and tablet dosages form. The method was further validated as per ICH guidelines (ICH Q2(R1), 2005) for the parameter like precision, accuracy, sensitivity, and linearity. The result of analysis was validated statistically and by recovery studies.

MATERIAL AND METHODS

Samples

ATC and FEN was kindly provided by Rightaid Laboratories, Hyderabad. The pharmaceutical formulation Mactor F used in this study was procured from local market of Bareilly.

Reagents

Methanol (GR grade), Hydrochloric acid (GR grade), Sodium hydroxide were obtained from Qualingens fine chemicals, Mumbai.

Instruments

UV-Visible double beam spectrophotometer (UV-3200 LAB INDIA) with 1 cm matched quartz cells, Digital balance (K-Roy Electronic), Oven (CLE-101, coslab) and volumetric flask, micropipette were utilized for present work.

Preparation of standard stock solution

An accurately weighed quantity of about 50 mg of ATC was taken in 50 ml volumetric flask and dissolved in sufficient quantity of methanol followed by sonication for 10 min and finally diluted to 50 ml with the same solvent so as to get the concentration of 1000 $\mu\text{g/ml}$. An accurately weighed quantity of about 50 mg of FEN was taken in 50 ml volumetric flask, dissolved in sufficient quantity of methanol and then sonicated for 15 min and finally diluted up to the mark with same solvent so as to get the concentration of 1000 $\mu\text{g/ml}$. From this solution, 5 ml was pipetted out in a 50 ml volumetric and volume was made up with methanol and get concentration 100 $\mu\text{g/ml}$ used for making dilution for calibration curve.

Determination of λ_{max}

The standard solution of ATC and FEN were separately scanned at different concentration in

the range of 200-400 nm and the λ_{\max} was determined for each drug.

Preparation of calibration curve

For each drug, appropriate aliquots were pipetted out from standard stock solution into the series of 10 ml volumetric flask and the volume was made up to the mark with methanol to get concentration of 6-16 $\mu\text{g/ml}$ of ATC and 2-12 $\mu\text{g/ml}$ of FEN. Solutions of different concentrations for each drug were analyzed at their respective wavelengths and absorbances were recorded.

Simultaneous equation method

Two wavelengths selected for the method (247 nm and 287 nm) that are absorption maximas of ATC and FEN respectively in methanol. Standard stock solution of 100 $\mu\text{g/ml}$ both the drug was prepared separately in methanol. The stock solution of both drug was further diluted separately with methanol to get series of standard solution of 6-16 $\mu\text{g/ml}$ for ATC and 2-12 $\mu\text{g/ml}$ for FEN. The absorbances were measured at the selected wavelengths and absorptivities (A 1%, 1 cm) for both the drugs at both wavelengths were determined as mean of three independent determinations. Concentrations in the sample were obtained by using following equations.

$$C_X = \frac{A_1 a_{y2} - A_2 a_{y1}}{a_{x1} a_{y2} - a_{x2} a_{y1}} \quad \text{Eq. 1}$$

$$C_Y = \frac{A_1 a_{x2} - A_2 a_{x1}}{a_{y1} a_{x2} - a_{y2} a_{x1}} \quad \text{Eq. 2}$$

where A_1 and A_2 are absorbances of mixture at 247 nm and 287 nm respectively, a_{x1} and a_{x2} are absorptivities of ATC at λ_1 and λ_2 respectively and a_{y1} and a_{y2} are absorptivities of FEN at λ_1 and λ_2 respectively. C_x and C_y are concentrations of ATC and FEN respectively.

Preparation for tablets assay

Twenty ATC and FEN tablets (10 mg atorvastatin and 160 mg fenofibrate) were weighed and powdered. A portion equivalent to 160 mg of fenofibrate was weighed into 100 ml clean and dry volumetric flask, About 70 ml of methanol was added and sonicated for 20 min; volume was made upto the mark with methanol, mixed well and filtered through Whatman filter paper No. 41. First few ml of filtrate was discarded and then 5 ml of filtrate was pipetted out and diluted to 50 ml with methanol. Then, the absorbances were recorded at the respective wavelengths.

Recovery study

To check the accuracy of the developed method recovery study was carried out as per ICH norms; where to a reanalyzed sample solution, standard solutions of both the drugs were added equivalent to 80, 100 and 120% of its drug content.

METHOD VALIDATION

The analytical method was validated with respect to parameters such as linearity, limit of detection (LOD), limit of quantitation (LOQ), precision, accuracy, robustness, and recovery (ICH Q2(R1), 2005).

Linearity

Linearity was established by least squares linear regression analysis of the calibration curve.

Accuracy

Accuracy was studied by adding two different amounts (corresponding to 80%, 100% and 120% of the test preparation concentrations) of ATR and FEN to the placebo preparation and comparing the actual and measured concentrations. For each level, three solutions were prepared and each was injected in duplicate.

Precision

The precision of the method, as intra-day repeatability was evaluated by performing six independent assays of the test sample preparation and calculating the RSD %. The intermediate (interday) precision of the method was checked by performing same procedure on different days by another person under the same experimental conditions.

LOD and LOQ

The LOD and LOQ of ATC and FEN were calculated by mathematical equation:

$$\text{LOD} = 3.3 \times \text{standard deviation} / \text{slope}$$

$$\text{LOQ} = 10 \times \text{standard deviation} / \text{slope}$$

Robustness

Robustness of proposed method was performed by changing UV analyst and remaining conditions (solvent, dilution, UV Spectrophotometer) were same.

FORCED DEGRADATION STUDIES

Forced degradation studies were performed on ATC and FEN to prove the stability indicating

property of the method. The stress conditions employed for degradation study included light exposure, acid hydrolysis (0.1 N HCl), base hydrolysis (0.1 N NaOH), thermal hydrolysis. The duration of time selected for degradation studies was 6 h (ICH Q1A(R2), 2003).

Acid hydrolysis

Solutions for acid degradation studies were prepared in methanol (12 $\mu\text{g/ml}$ for ATC and 8 $\mu\text{g/ml}$ for FEN) and added 10 ml 0.1 M hydrochloric acid solution and kept at room temperature (22°C). It was observed that both acid and base hydrolysis were fast reactions for both drugs and almost completed within 3 h of the sample preparation, therefore the samples were analyzed at 247 nm for ATC and 287 nm for FEN after this period of time.

Base hydrolysis

Solutions for base degradation studies were prepared in methanol (12 $\mu\text{g/ml}$ for ATC and 8 $\mu\text{g/ml}$ for FEN) and 100 ml 0.1 M sodium hydroxide add in both dilution and kept at room

temperature (22°C) and the resultant solutions analyzed 10 min after preparation at 247 nm for ATC and 287 nm for FEN.

Photostability studies

Fifty mg drug was weighed and kept in the sun light for 12 h, after that the solutions for photostability studies were prepared in methanol and the dilutions (12 $\mu\text{g/ml}$ - ATC, 8 $\mu\text{g/ml}$ - FEN) were prepared & analyzed in UV spectrophotometer at 247, 287 nm for ATC, FEN.

Thermal degradation

Fifty mg drug was weighed and kept in the oven and temperature was maintained at 80°C for 3 h, after that the solutions for photostability studies were prepared in methanol and the dilution (12 $\mu\text{g/ml}$ - ATC and 8 $\mu\text{g/ml}$ - FEN) were prepared and analyzed in UV spectrophotometer.

RESULTS AND DISCUSSION

The UV scanning showed spectrum exhibiting λ_{max} of 247 nm and 287 nm for ATC and FEN respectively (**Figure 2**).




Fig. 2. λ_{max} of ATC and FEN

The linearity of the proposed method was investigated in the range of 6-16 $\mu\text{g/ml}$ and 2-12 $\mu\text{g/ml}$ for ATC, FEN respectively. Calibration curves showed a linear relationship between the

absorbance and concentration. The line equation for ATC ($y = 0.041x + 0.043$) with r^2 of 0.999 and for FEN ($y = 0.054x - 0.003$) with r^2 of 0.999 was obtained (**Table 1, Figure 3**).

Table 1. Calibration curve parameters

S. No.	Parameter	Atorvastatin calcium	Fenofibrate
1	Linearity range ($\mu\text{g/ml}$)	6-16 $\mu\text{g/ml}$	2-12 $\mu\text{g/ml}$
2	Correlation coefficient (r^2)	0.999	0.999
3	Slope	0.041	0.054
4	Intercept	0.043	0.003




Fig. 3. Calibration curve of Atorvastatin and Fenofibrate

The results of tablet assay analysis was found within the prescribed limits and complied with

pharmacopoeial standards. The results of tablet analysis are summarized in **Table 2**.

Table 2. Analysis of tablet dosage form

Formulation	Drug	Label claim (mg)	% Label claim (Mean \pm SD)
Tablet	ATC	10 mg	101.33 \pm 0.0018
	FEN	160 mg	104.12 \pm 0.0034

The overlaid spectra of ATC and FEN exhibited λ_{\max} of 247 nm and 287 nm for ATC and FEN

respectively which were quite clearly separated from each other (**Figure 4**).




Fig. 4. Overlaid spectra of ATC and FEN

The constructed calibration curve was linear over the constructed range for fenofibrate (**Table 3**). The LOD of ATC and FEN were found to be 0.2695 $\mu\text{g/ml}$ and 0.0222 $\mu\text{g/ml}$ and the LOQ of ATC and FEN were found to be 0.8780 $\mu\text{g/ml}$ and 0.22221 $\mu\text{g/ml}$. Validation was performed as per ICH guidelines (ICH Q2(R1), 2005) for Linearity, accuracy, precision, LOD and LOQ. The LOD-0.2695 $\mu\text{g/ml}$, 0.0222 $\mu\text{g/ml}$ for ATC and FEM and the LOQ 0.8780 $\mu\text{g/ml}$, 0.222

$\mu\text{g/ml}$ for ATC and FEM respectively. The results of method validation parameters are summarized in **Table 4**. The result of forced degradation are compiled in **Table 5**. The stability studies of ATC and FEN were conducted and the degradation characteristics were found to be much more prominent in acid hydrolysis in FEN and alkaline hydrolysis in ATC (**Figure 5, 6**). The spectrum of photo and thermal degradation studies are evident in **Figure 7, 8**.

Table 3. Linearity study of FEN at 287 nm in methanol

Conc. ($\mu\text{g/ml}$)	Dilution 1	Dilution 2	Dilution 3	Dilution 4	Dilution 5	Dilution 6	Mean \pm SD*
2	0.1118	0.1072	0.1085	0.1060	0.1058	0.1079	0.1078 \pm 0.0020
4	0.2167	0.2185	0.2177	0.2171	0.2161	0.2166	0.2170 \pm 0.00086
6	0.3222	0.3234	0.3243	0.3227	0.3241	0.3231	0.3233 \pm 0.00073
8	0.4330	0.4324	0.4342	0.4339	0.4333	0.4337	0.4334 \pm 0.00059
10	0.5478	0.5490	0.5495	0.5467	0.5497	0.5438	0.5477 \pm 0.0020
12	0.6543	0.6556	0.6564	0.6550	0.6555	0.6584	0.6558 \pm 0.001298

*SD - Standard deviation

Table 4. Validation parameter for ATC and FEN

S. No.	Parameter (units)	ATC	FEN
1.	Linearity	6-16 $\mu\text{g/ml}$	2-12 $\mu\text{g/ml}$
2.	Accuracy (80%)	99.38 \pm 0.0009	99.30 \pm 0.0014
	Accuracy (100%)	98.52 \pm 0.0013	98.69 \pm 0.0008
	Accuracy (120%)	98.32 \pm 0.0010	97.34 \pm 0.0029
3.	Interday Precision (1 st day)	109.87% \pm 0.0034*	99.21% \pm 0.00216
	(2 nd day)	109.66% \pm 0.0123*	100.09% \pm 0.0014
	(3 rd day)	123.38% \pm 0.0013*	106.10% \pm 0.0016
4.	Intraday Precision 1 st h	109.87% \pm 0.0034*	99.21% \pm 0.0026
	2 nd h	106.22% \pm 0.009*	100.96% \pm 0.0001
	3 rd h	100.44% \pm 0.0013*	99.88% \pm 0.002
5.	LOD	0.2695 ($\mu\text{g/ml}$)	0.0222 ($\mu\text{g/ml}$)
6.	LOQ	0.8780 ($\mu\text{g/ml}$)	0.2222 ($\mu\text{g/ml}$)
7.	Robustness	95.81% \pm 0.0007*	105% \pm 0.0008

*Standard deviation

Table 5. Results of forced degradation study

S. No.	Condition	Absorbances (λ)		Mean \pm SD*		Result (% degradation)	
		ATC	FEN	ATC	FEN	ATC	FEN
1.	Acid degradation	255 nm	No absorbance	0.2720 \pm 0.0021	0.4128 \pm 0.0009	39.12%	100%
2.	Alkaline degradation	249 nm	301 nm	0.2936 \pm 0.0043	0.3025 \pm 0.0125	34.29%	30.25%
3.	Thermal degradation	247 nm	286.5 nm	0.1993 \pm 0.0002	0.3645 \pm 0.0028	55.4%	100%
4.	Photolytic degradation	247 nm	287 nm	0.3874 \pm 0.0109	0.6977 \pm 0.0022	13.3%	20.0%

*SD - Standard deviation




Fig. 5. Acid degradation of ATC and FEN




Fig. 6. Base degradation of ATC and FEN




Fig. 7. Photostability degradation of ACE and FEN




Fig. 8. Thermal degradation of ATC and FEN

CONCLUSION

The proposed method is found to be simple, sensitive and reproducible and hence it can be used in routine analysis for simultaneous determination of stability indicating UV spectrophotometric method for ATC and FEN in bulk as well as in pharmaceutical preparation. Statistical analysis of the results has been carried

REFERENCES

- Altuntas TG, Erk N, Liquid chromatographic determination of atorvastatin in bulk drug, tablets and human plasma. *J. Liq. Chromatogr. Relat. Technol.* 2004;27(1):83-93. [DOI: 10.1081/JLC-120027087]
- Diwan VP, Anusha N, Huidrom S, Surendra N, Prasad VVLN. Stability indicating high performance liquid chromatographic assay for the simultaneous determination of atorvastatin calcium and fenofibrate in tablet dosage form. *Int. J. Pharm. Ind. Res.* 2012;2(3):331-9.
- Erk N. Extractive spectrophotometric determination of atorvastatin in bulk and pharmaceutical formulations. *Anal. Lett.* 2003;36(12):2699-711. [DOI: 10.1081/AL-120024645]
- Erturk S, Aktas ES, Ersoy L, Ficioglu S. An HPLC method for the determination of atorvastatin and its impurities in bulk drugs and tablets. *J. Pharm. Biomed. Anal.* 2003;33(5):1017-23. [DOI: 10.1016/S0731-7085(03)00408-4]
- Gupta KR, Askarkar SS, Rathod PR, Wadodkar SG. Validated spectrophotometric determination of fenofibrate in formulation, *Der Pharm. Sin.* 2010;1(1):173-8.
- ICH, Q1A(R2), Stability testing of new drug substances and products, International Conference on Harmonization, IFPMA, Geneva, 2003.
- ICH, Q2(R1), Validation of analytical procedures: Text and

out revealing high accuracy and good precision.

ACKNOWLEDGEMENT

Authors are thankful to project guide and management, Invertis Institute of Pharmacy, Invertis University, Bareilly, Uttar Pradesh for providing equipment and facility during entire duration of project work.

- methodology, International Conference on Harmonization, IFPMA, Geneva, 2005.
- Indian Pharmacopoeia, Indian Pharmacopoeia Commission, Ghaziabad, 2007;2:131-3.
- Janardhanan VS, Manavalan R, Valliappan K. Stability-indicating HPLC method for the simultaneous determination of pantoprazole, rabeprazole, lansoprazole and domperidone from their combination dosage forms. *Int. J. Drug Dev. Res.* 2011;3(4):323-35.
- Salama FMM, Nassar MWI, Sharaf El-Din MMK, Attia KAM, Kaddah MY, Determination of fenofibrate and the degradation product using simultaneous UV-derivative spectrometric method and HPLC. *Am. J. Anal. Chem.* 2011;2(3):332-43. [DOI: 10.4236/ajac.2011.23042]
- Singh S, Dubey N, Jain DK. Simultaneous estimation of atorvastatin, clopidogrel and aspirin in capsule dosage form using UV-spectroscopy. *Asian J. Res. Chem.* 2010; 3(4):885-7.
- Singh S, Yadav AK, Gautam H. Simultaneous estimation of valsartan and hydrochlorothiazide in solid dosage form using UV spectroscopy. *Bull. Pharm. Res.* 2011;1(3):10-2.
- Singh S, Yadav AK, Gautam H. First order derivative spectrophotometric determination of telmisartan in pharmaceutical formulation. *Bull. Pharm. Res.* 2012;2(2): 83-6.
- <http://www.chemspider.com/Chemical-Structure.3222.html>
