



RESEARCH ARTICLE

SYNTHETIC STUDIES ON NOVEL NITROQUINAZOLINONE ANALOGS WITH ANTIMICROBIAL POTENTIAL

Rajiv Dahiya^{1*} and Rita Mourya²

¹Department of Pharmaceutical Chemistry, Globus College of Pharmacy, Bhopal-462 045, Madhya Pradesh, India

²Research Scholar, Institute of Pharmaceutical Science and Research Center, Bhagwant University, Sikar Road, Ajmer-305 004, Rajasthan, India

*E-mails: drrajivdahiya@gmail.com, ritz_pharma@yahoo.co.in

Tel.: +91 9630229885, +91 8878603338.

Received: March 16, 2013 / Revised: May 20, 2013 / Accepted: May 21, 2013

A novel series of 4-[2-(3-bromophenyl)-7-nitro-4-oxo-3,4-dihydro-3-quinazolinyl]benzoyl amino acids and di/tripeptides was synthesized using diisopropylcarbodiimide (DIPC) as the coupling agent and *N*-methylmorpholine (NMM) as the base. Structures of all the newly synthesized peptide analogs were elucidated using IR, ¹H/¹³C NMR, MS spectral data and evaluated for antimicrobial potential against pathogenic microbes. Most of the compounds exhibited potent antifungal activity against pathogenic *Candida albicans* and dermatophytes, in comparison to reference compound. Good bioactivity was also seen against gram-negative bacteria for synthesized compounds.

Key words: Quinazolinones, 4-Nitroanthranilic acid, Peptide analogs, Coupling, Antifungal activity.


INTRODUCTION

Literature is enriched with several works on synthesis of potent quinazolinone and benzoic acid derivatives with diverse biological activities (de Moura *et al* 2004; Trusheva *et al* 2010; Wu *et al* 2010; Sharma *et al* 2011; Wu *et al* 2011) but very few reports have been yet received regarding peptide coupling of quinazolinones, although potent peptide analogs of aroylbenzoic acid, furoic acid, aryloxyacetic acid, coumarin, quinoxaline, benzothiophene, benzimidazole and imidazole are already reported (Poojary *et al* 2001; 2003; Himaja *et al* 2002; 2003; Dahiya and Pathak, 2006a; 2006b; 2006c; Dahiya *et al* 2006a; 2006b; Dahiya and Pathak, 2007; Dahiya and Kaur, 2007a; 2007b; 2008; Dahiya, 2008a, 2008b; Dahiya *et al* 2008a; 2008b; Dahiya and Kumar, 2008; Dahiya and Bansal, 2008; Dahiya *et al* 2010). Keeping in view the pharmacological potential of quinazolinone and benzoic acid derivatives, both moieties were coupled in single nucleus and further, in continuation of our work on synthesizing potent antimicrobial peptide analogs (Dahiya and Mourya, 2012), a novel series of 4-[2-(3-bromophenyl)-7-nitro-4-oxo-

3,4-dihydro-3-quinazolinyl]benzoyl amino acids and peptides was synthesized to get bioactive agents with biological interest. 4-[2-(3-bromophenyl)-7-nitro-4-oxo-3,4-dihydro-3-quinazolinyl]benzoic acid (**1**) was prepared by interaction of *p*-aminobenzoic acid and 2-(3-bromophenyl)-7-nitro-4*H*-3,1-benzoxazin-4-one, which was in turn prepared from the 4-nitroanthranilic acid and 3-bromobenzoyl chloride at 0-5 °C in presence of pyridine (Gao *et al* 2007). Dipeptides Boc-Pro-Pro-OMe, Boc-Try-His-OMe, Boc-His-Phe-OMe, were prepared from the corresponding Boc-amino acids and amino acid methyl ester hydrochlorides using the dicyclohexylcarbodiimide (DCC) and triethylamine (TEA) in dichloromethane (DCM). Similarly, Boc-Phe-Ile-Pro-OMe and Boc-His-Tyr-His-OMe was prepared by coupling Boc-Phe/Boc-His with Ile-Pro-OMe/Tyr-His-OMe in alkaline conditions. For coupling, di/tripeptide units were selected from pharmacologically active natural as well as synthetic cyclic polypeptides (Dahiya and Pathak, 2006d; Dahiya and Gautam, 2011).

The free acid **1** was coupled with various L-amino acid methyl ester hydrochlorides using diisopropylcarbodiimide (DIPC) as the coupling agent to get newly synthesized 4-[2-(3-bromo phenyl)-7-nitro-4-oxo-3,4-dihydro-3-quinazolin-yl]benzoyl amino acid methyl ester (**1a**).

Similarly, dipeptide and tripeptide coupling was done to get dipeptide (**1b-d**) and tripeptide (**1e, 1f**) methyl ester derivatives. Finally, selected compounds were hydrolyzed with lithium hydroxide to get (**1g-j**), corresponding acid derivatives (**Scheme 1**).



X = Try (**1a, 1g**), Pro-Pro (**1b**), Try-His (**1c, 1h**), His-Phe (**1d, 1i**), Phe-Ile-Pro (**1e**), His-Tyr-His (**1f, 1j**)

a = pyridine, 0 °C; b = PABA, 170 °C, 2h; c = DIPC, NMM, DCM, RT, 24 h; d = LiOH, THF:H₂O (1:1), RT, 1 h

Scheme 1. Synthetic pathway for novel quinazolinopeptide analogs (**1a-j**)

All the compounds were synthesized in good yields and their structures were confirmed by FTIR, ¹H/¹³C NMR and Mass spectra. In addition,

elemental analysis of the newly synthesized compounds was performed for carbon, hydrogen and nitrogen content (**Table 1**).

Table 1. Physical data of newly synthesized quinazolinopeptide analogs

Compd	X	M.P. (°C)	Yield (%)	C (%) Calcd./Found	H (%) Calcd./Found	N (%) Calcd./Found
1a	L-Try	122	78	59.47/59.49	3.63/3.59	10.51/10.52
1b	L-Pro-L-Pro	–	89	56.98/56.99	4.18/4.15	10.38/10.40
1c	L-Try-L-His	188	79	58.29/58.27	3.89/3.92	13.94/13.95
1d	L-His-L-Phe	207	80	58.12/58.09	3.95/3.96	12.82/12.85
1e	L-Phe-L-Ile-L-Pro	166	95	60.22/60.19	4.93/4.95	10.03/10.05
1f	L-His-L-Tyr-L-His	223	92	56.28/56.25	4.06/4.08	15.26/15.25
1g	L-Try	159	70	58.91/58.89	3.40/3.43	10.73/10.75
1h	L-Try-L-His	145	67	57.80/57.78	3.70/3.73	14.19/14.20
1i	L-His-L-Phe	192	71	57.61/57.59	3.76/3.75	13.06/13.09
1j	L-His-L-Tyr-L-His	179	89	55.82/55.79	3.90/3.92	8.84/8.85

EXPERIMENTAL**Materials and Methods**

Melting points were determined by open capillary method and are uncorrected. L-Amino acids, di-*tert*-butylpyrocarbonate (Boc₂O), diisopropylcarbodiimide (DIPC), trifluoroacetic acid (TFA), *N*-methylmorpholine (NMM) and triethylamine (TEA) were obtained from Spectrochem Limited, Mumbai, India. IR spectra were recorded on fourier transform infrared spectrophotometer (Jasco, Japan) using KBr pellets/CHCl₃ for all the synthesized compounds. ¹H (300 MHz) and ¹³C (75 MHz) NMR spectra were recorded on Bruker spectropin spectrometer using CDCl₃ as solvent and TMS as an internal standard. Mass spectra were recorded on Micromass Quattro II triple quadrupole mass spectrometer at 70 eV. Elemental analysis of all compounds were performed on CHN Analyzer (Elementar, Germany). Purity of all the compounds was checked by TLC on precoated silica gel G plates.

Preparation of peptides

Peptide units were prepared by stirring a mixture of amino acid methyl ester hydrochloride and Boc-amino acid/dipeptide (0.01 mol each) with DCC (0.01 mol), TEA (0.021 mol), and DCM (50 ml) for 24 h. Then, reaction mixture was filtered and the residue was washed with DCM (25 ml). After washing the filtrate with 5% NaHCO₃ and saturated NaCl solutions, organic layer was dried over anhyd. Na₂SO₄ and evaporated in vacuum after filtration. The crude product was recrystallized from a mixture of chloroform and petroleum ether followed by cooling at 0 °C. Resulting Boc-di/tripeptide methyl ester (0.01 mol) was dissolved in CHCl₃ (25 ml) and treated with trifluoroacetic acid (0.02 mol). The mixture was stirred at RT for 1 h and washed with saturated NaHCO₃ solution. The resulting organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. Finally, crude product was purified by recrystallization with CHCl₃ and petroleum ether to get the deprotected di/tripeptide methyl ester. For protecting L-amino acids at amino end, Boc₂O was used whereas the carboxyl group of L-amino acids was protected by esterification with methanol using thionyl chloride. Peptides were prepared by Bodanzky method with certain modifications (Bodanzsky and Bodanzsky, 1984). Furthermore, trifluoroacetic acid was used for the removal of Boc group and ester group was

removed by alkaline hydrolysis with lithium hydroxide.

Preparation of 4-[2-(3-bromophenyl)-7-nitro-4-oxo-3,4-dihydro-3-quinazolinyl]-benzoic acid (1)

To a cold solution of 4-nitroanthranilic acid (0.01 mol) in pyridine (30 ml) maintained at 0°C, another solution of 3-bromobenzoyl chloride (0.02 mol) in pyridine (30 ml) was added slowly with constant stirring. The reaction mixture was further stirred for 30 min at RT and set aside for 1 h. The pasty mass obtained was diluted with water (50 ml) and treated with aqueous sodium bicarbonate solution. The precipitate obtained was filtered off, washed with water and finally dried. Equimolar amounts (0.025 mol) of precipitate and *p*-aminobenzoic acid (PABA) were heated at 170-180 °C for 2 h in an oil bath. The separated jelly like mass solidified upon cooling. The crude product was finally recrystallized from ethanol to give a good yield of title compound as pale yellow solid; yield 78%; M.P. 193 °C; IR (KBr): 3298-2509 (O-H str, COOH), 3078-3069, 3054-3049 (C-H str, rings), 1706 (C=O str, COOH), 1667 (C=O str, ring), 1596 (C=N str, ring), 1584-1575, 1422-1414 (C=C str, rings), 1515, 1350 (NO₂ str, asym and sym), 1402 (O-H def, COOH), 863 (C-N str, Ar-NO₂), 779, 735, 721, 702, 698, 663 (C-H def, oop, rings), 685, 622 (C-Br str) cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 10.76 (1H, s, OH, COOH), 10.76 (1H, s, H-8, quinazolinone ring [qze]), 8.34 (1H, d, J = 7.4 Hz, H-6, qze), 8.28 (1H, d, J = 7.6 Hz, H-5, qze), 8.15 (1H, s, H-2, bromobenzene moiety [bbz]), 7.91-7.89 (1H, d, J = 6.9 Hz, H-2, bbz), 7.85-7.83 (2H, dd, J = 6.95 Hz, 7.2 Hz, m-H's, benzoic acid moiety [bza]), 7.42-7.38 (3H, m, o-H's, bza and H-4, bbz), 7.42-7.38 (1H, t, J = 6.6 Hz, H-5, bbz) ppm; ¹³C NMR (CDCl₃, 70 MHz): δ 171.5 (C=O, -COOH), 168.2 (C-2, qze), 157.9 (C-4, qze), 154.6 (C-7, qze), 149.2 (C-2', qze), 138.6 (C-1, bza), 133.7 (C-1, bbz), 133.2 (2C, m-C's, bza), 132.6 (2C, o-C's, bza), 130.4 (C-4, bbz), 129.9 (p-C, bza), 129.2 (C-2, bbz), 128.8 (C-5, bbz), 128.2 (C-5, bbz), 127.7 (C-5, qze), 125.1 (C-3', qze), 123.4 (C-3, bbz), 118.8 (C-6, qze), 116.5 (C-8, qze) ppm.

General procedure for preparation of 4-[2-(3-bromophenyl)-7-nitro-4-oxo-3,4-dihydro-3-quinazolinyl]benzoyl amino acid and peptide derivatives (1a-j)

Title compounds 1a-j were prepared by the coupling of 4-[2-(3-bromophenyl)-7-nitro-4-

oxo-3,4-dihydro-3-quinazoliny]benzoic acid (**1**) with amino acid methyl ester hydrochlorides/peptide methyl esters by using DIPC and NMM in DCM by following the same procedure as that adopted for peptide synthesis to get the amino acid and peptide derivatives (**1a-j**). Amino acid (**1a**), dipeptide (**1c**, **1d**) and tripeptide (**1f**) derivatives were further hydrolyzed with LiOH to get the corresponding free acids (**1g-j**).

For removal of the ester group, peptide ester derivative (0.01 mol) was dissolved in THF:H₂O (1:1) and LiOH (0.015 mol) was added to the solution at 0 °C. The resulting mixture was stirred at RT for 1 h and then acidified to pH 3.5 with 1 N H₂SO₄. The aqueous layer was extracted with Et₂O (2 × 20 ml) and the combined organic extracts were dried over Na₂SO₄ and concentrated under reduced pressure to get the desired hydrolyzed compound.

1a (C₃₃H₂₄BrN₅O₆): ¹³C NMR (CDCl₃, 70 MHz): δ 172.3 (C=O, ester), 169.0 (C-2, qze), 168.6 (C=O, bza), 158.3 (C-4, qze), 154.1 (C-7, qze), 148.9 (C-2', qze), 139.7 (C-1, bza), 137.8 (C-2', indole ring [idr]), 134.2 (C-1, bbz), 131.9 (C-4, bbz), 130.8 (C-2, bbz), 129.2 (C-5, bbz), 128.7 (C-6, bbz), 128.0 (C-5, qze), 127.5 (2C, m-C's, bza), 127.0 (2C, o-C's, bza), 126.8 (C-3', idr), 125.6 (p-C, bza), 124.8 (C-3', qze), 123.1 (C-3, bbz), 122.6 (C-2, idr), 121.7 (C-6, idr), 119.5 (C-5, idr), 118.7 (C-4, idr), 117.9 (C-6, qze), 116.6 (C-8, qze), 114.4 (C-3, idr), 112.9 (C-7, idr), 55.6 (C-α, Trp), 51.7 (OCH₃), 19.9 (C-β, Trp) ppm.

1b (C₃₂H₂₈BrN₅O₇): ¹H NMR (CDCl₃, 300 MHz): δ 8.64 (1H, s, H-8, qze), 8.44-8.39 (2H, m, m-H's, bza), 8.31-8.26 (2H, m, H-5 and H-6, qze), 8.11 (1H, s, H-2, bbz), 7.92-7.89 (1H, t, J = 6.5 Hz, H-6, bbz), 7.62-7.58 (2H, m, o-H's, bza), 7.40-7.38 (1H, d, J = 6.85 Hz, H-4, bbz), 7.29-7.26 (1H, t, J = 6.7 Hz, H-5, bbz), 4.41-4.38 (1H, t, J = 6.9 Hz, H-α, Pro-1), 4.26-4.23 (1H, t, J = 6.85 Hz, H-α, Pro-2), 3.72-3.69 (2H, t, J = 7.15 Hz, H-δ, Pro-2), 3.62 (3H, s, OCH₃), 3.45-3.42 (2H, t, J = 7.2 Hz, H-δ, Pro-1), 2.69-2.64 (2H, m, H-β, Pro-1), 2.05-2.01 (2H, m, H-β, Pro-2), 1.97-1.92 (4H, m, H-γ, Pro-1 and Pro-2) ppm.

1c (C₃₉H₃₁BrN₈O₇): IR (KBr): 3486-3479 (N-H str, heterocyclic rings), 3129, 3122 (N-H str, amide), 3085-3069, 3055, 3047-3042 (C-H str, rings), 2925 (C-H str, asym, CH₂), 2844 (C-H str, sym, CH₂), 1745 (C=O str, ester), 1664 (C=O str, ring), 1639-1634 (C=O str, 2° amide), 1599-1595

(C=N str, rings), 1594-1573, 1438-1423 (C=C str, rings), 1537, 1532 (N-H bend, 2° amide), 1515, 1357 (NO₂ str, asym and sym), 1274 (C-O str, ester), 864 (C-N str, Ar-NO₂), 854-843, 775, 729-714, 692, 657-653 (C-H def, oop, rings), 681, 622 (C-Br str) cm⁻¹.

1d (C₃₇H₃₄BrN₅O₇): ¹³C NMR (CDCl₃, 70 MHz): δ 171.6 (C=O, His), 170.3 (C=O, ester), 169.4 (C-2, qze), 167.7 (C=O, bza), 158.0 (C-4, qze), 152.3 (C-7, qze), 149.1 (C-2', qze), 142.8 (C-2, imidazole ring [imz]), 140.2 (C-1, bza), 137.3 (C-γ, Phe), 134.7 (C-1, bbz), 133.5 (C-4, bbz), 132.4 (C-2, bbz), 130.2 (C-5, bbz), 129.4 (2C, o-C's, Phe), 128.9 (C-6, bbz), 128.3 (2C, m-C's, Phe), 128.1 (2C, o-C's, bza), 127.9 (C-5, qze), 127.5 (2C, m-C's, bza), 126.7 (p-C, Phe), 125.6 (C-4, imz), 125.0 (p-C, bza), 124.5 (C-3', qze), 123.6 (C-3, bbz), 120.5 (C-6, qze), 118.8 (C-8, qze), 116.6 (C-5, imz), 59.2 (C-α, His), 55.8 (C-α, Phe), 52.7 (OCH₃), 38.6 (C-β, Phe), 18.8 (C-β, His) ppm.

1e (C₄₂H₄₁BrN₆O₈): MS *m/z* (rel. int.): 14 (4.7), 15 (14.2), 29 (9.8), 31 (9.6), 59 (12.4), 70 (11.5), 77 (16.4), 86 (17.1), 91 (21.5), 120 (13.9), 145 (11.8), 156 (11.4), 190 (16.9), 345 (24.5), 421 (32.6), 449 (68.2), 568 (10.7), 596 (100), 681 (24.7), 709 (74.7), 778 (16.6), 806 (24.2), 822 (19.4), 837 (M⁺, 2.9), 838 (M⁺, 0.6).

1f (C₄₃H₃₇BrN₁₀O₉): MS *m/z* (rel. int.): 14 (3.9), 15 (3.6), 17 (4.7), 31 (9.9), 59 (13.3), 67 (15.7), 81 (22.9), 93 (9.4), 107 (11.8), 110 (19.5), 136 (14.8), 145 (12.6), 156 (10.9), 190 (16.9), 345 (21.2), 421 (29.4), 449 (65.6), 558 (14.3), 586 (100), 721 (14.6), 749 (70.2), 858 (15.9), 886 (33.7), 902 (17.2), 917 (M⁺, 2.6), 918 (M⁺, 0.8).

1g (C₃₂H₂₂BrN₅O₆): ¹³C NMR (CDCl₃, 70 MHz): δ 175.8 (C=O, -COOH), 169.1 (C=O, bza), 168.6 (C-2, qze), 157.8 (C-4, qze), 152.9 (C-7, qze), 148.0 (C-2', qze), 140.5 (C-1, bza), 136.9 (C-2', idr), 133.7 (C-1, bbz), 132.3 (C-4, bbz), 130.4 (C-2, bbz), 130.1 (C-6, bbz), 129.6 (C-5, bbz), 128.4 (C-5, qze), 128.0 (2C, m-C's, bza), 127.6 (2C, o-C's, bza), 127.1 (C-3', idr), 125.4 (C-3', qze), 124.5 (p-C, bza), 123.5 (C-2, idr), 123.0 (C-3, bbz), 122.2 (C-6, idr), 119.2 (C-5, idr), 118.2 (C-6, qze), 117.7 (C-4, idr), 117.2 (C-8, qze), 115.4 (C-3, idr), 111.5 (C-7, idr), 54.9 (C-α, Trp), 19.4 (C-β, Trp) ppm.

1h (C₃₈H₂₉BrN₈O₇): IR (KBr): 3488-3481 (N-H str, heterocyclic rings), : 3298-2487 (O-H str, -COOH), 3127, 3120 (N-H str, amide), 3087-

3072, 3058, 3044-3036 (C-H str, rings), 2928 (C-H str, asym, CH₂), 2845 (C-H str, sym, CH₂), 1716 (C=O str, COOH), 1668 (C=O str, ring), 1639-1632 (C=O str, 2° amide), 1596-1592 (C=N str, rings), 1597-1577, 1439-1425 (C=C str, rings), 1535, 1530 (N-H bend, 2° amide), 1516, 1359 (NO₂ str, asym and sym), 868 (C-N str, Ar-NO₂), 856-842, 777, 726-717, 698, 655-649 (C-H def, oop, rings), 685, 626 (C-Br str) cm⁻¹.

1i (C₃₆H₂₈BrN₇O₇): ¹³C NMR (CDCl₃, 70 MHz): δ 173.5 (C=O, COOH), 171.4 (C=O, His), 169.2 (C-2, qze), 168.3 (C=O, bza), 158.6 (C-4, qze), 152.0 (C-7, qze), 148.8 (C-2', qze), 141.9 (C-2, imz), 138.1 (C-γ, Phe), 137.7 (C-1, bza), 134.2 (C-1, bbz), 133.9 (C-4, bbz), 132.0 (C-2, bbz), 130.6 (C-5, bbz), 129.7 (2C, o-C's, Phe), 128.8 (C-5, qze), 128.5 (2C, o-C's, bza), 128.3 (C-6, bbz), 127.9 (2C, m-C's, bza), 127.5 (2C, m-C's, Phe), 126.3 (C-4, imz), 125.2 (p-C, bza), 124.8 (C-3', qze), 124.5 (p-C, Phe), 124.0 (C-3, bbz), 120.9 (C-6, qze), 119.0 (C-8, qze), 115.8 (C-5, imz), 58.9 (C-α, His), 54.5 (C-α, Phe), 37.8 (C-β, Phe), 18.9 (C-β, His) ppm.

1j (C₄₂H₃₅BrN₁₀O₉): MS *m/z* (rel. int.): 14 (3.5), 17 (8.2), 45 (10.8), 67 (15.4), 81 (22.2), 93 (9.8), 107 (12.5), 110 (20.2), 136 (14.5), 145 (12.9), 156 (11.4), 190 (16.2), 345 (20.6), 421 (27.9), 449 (65.2), 558 (14.2), 586 (100), 721 (14.9),

749 (69.7), 858 (15.5), 886 (32.9), 903 (M⁺, 3.8), 904 (M⁺, 0.5).

Antimicrobial activity

Modified Kirby-Bauer disc diffusion method (Bauer *et al* 1966) was utilized for the testing antibacterial activity against four pathogenic bacteria *Bacillus subtilis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Escherichia coli*. MIC values of test compounds were determined by tube dilution technique. Ciprofloxacin was used as a reference compound and solvent (DMF) as negative control. Test sample and reference drug were tested at the concentration of 12.5-6 μg/ml. Serial plate dilution method (Khan, 1997) was employed for the evaluation of antifungal activity against three fungal strains, including dermatophytes *Microsporum audouinii*, *Trichophyton mentagrophytes* and diamorphic fungi *Candida albicans*. MIC values of test compounds were determined by employing the same technique as used for antibacterial studies using DMSO instead of DMF. During antifungal evaluation, test sample and reference drug – Griseofulvin were tested at the concentration of 6 μg/ml. The results of antibacterial and antifungal activity studies are compiled in **Table 2**. The detailed procedure for antimicrobial evaluation is reported in our previously published report (Dahiya and Pathak, 2007).

Table 2. Antimicrobial activity data of newly synthesized quinazolinopeptide analogs

Compound	Zone of inhibition (in mm)						
	<i>B.sub.</i>	<i>E.coli</i>	<i>S.aur.</i>	<i>P.aeru.</i>	<i>M.audo.</i>	<i>T.menta.</i>	<i>C.alb.</i>
1a	11	16	11	22	15	15	14
1b	14	-	13	20	-	16	16
1c	10	13	10	19	18	18	19
1d	10	13	10	19	15	17	16
1e	-	11	-	21	13	17	-
1f	9	13	11	19	17	19	19
1g	12	18	14	24	16	17	16
1h	11	14	11	21	20	20	21
1i	12	15	12	22	16	18	17
1j	11	14	14	21	18	21	22
Control	-	-	-	-	-	-	-
Ciprofloxacin	20	19	20	25	-	-	-
Griseofulvin	-	-	-	-	17	18	18

RESULTS AND DISCUSSION

All novel quinazolinopeptide analogs were synthesized successfully with good yield. All the synthesized compounds were found to exhibit

good antimicrobial activity against pathogenic dermatophytes, *Candida albicans* and gram-negative bacterium *Pseudomonas aeruginosa*. DIPC was found to be a good coupling agent both

economically as well as yield wise. IR spectra of synthesized compounds showed characteristic amide I and amide II bands of the -CO-NH- moiety and mass spectra showed molecular ion peak (M^+) at m/z values which were in consistent with the molecular formulas. Comparison of the antimicrobial activity data suggested that greater activity is observed in compounds with heterocyclic amino acid residues in their chain.

CONCLUSION

Almost, all the synthesized compounds were found to exhibit potent antifungal activity and moderate to good antibacterial activity against gram-negative bacteria. No significant bioactivity was observed against gram-positive bacteria for newly synthesized peptide analogs. Compounds (**1c**, **1f**) and their hydrolyzed

derivatives (**1h**, **1j**) were found to be more active than standard drug against pathogenic fungus *C. albicans* and dermatophytes whereas compounds (**1a**, **1g**) displayed good activity against bacteria *E. coli* and *P. aeruginosa*. Comparison of antimicrobial data has suggested that amino acid and peptide derivatives (**1g-j**) were more potent antifungal and antibacterial agents than corresponding methyl ester derivatives (**1a**, **1c**, **1d** and **1f**). On passing toxicity tests, these peptide derivatives may prove good candidate for clinical studies and can be new antifungal agents of tomorrow.

ACKNOWLEDGEMENT

Authors are thankful to Sophisticated Analytical Instrumental Laboratory (SAIL), RGPV, Bhopal (MP) and Jamia Hamdard University, New Delhi (India) for spectral and elemental analysis.

REFERENCES

- Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* 1966;45(4):493-6.
- Bodanzsky M, Bodanzsky A. The practice of peptide synthesis. Springer, New York: 1984; 68, 143, 207.
- Dahiya R. Synthesis, characterization and antimicrobial studies on some newer imidazole analogs. *Sci. Pharm.* 2008a;76(2):217-39. [DOI: 10.3797/scipharm.0803-04]
- Dahiya R. Synthesis of 4-(2-methyl-1H-5-imidazolyl) benzoyl amino acids and peptides as possible anthelmintic agents. *Ethiop. Pharm. J.* 2008b;26(1):17-26.
- Dahiya R, Bansal Y. Synthesis and antimicrobial potential of novel quinoxalinopeptide analogs. *Res. J. Chem. Environ.* 2008;12(3):52-8.
- Dahiya R, Gautam H. Solution phase synthesis and bioevaluation of cordyheptapeptide B. *Bull. Pharm. Res.* 2011;1(1):1-10.
- Dahiya R, Kaur R. Synthesis and biological screening of a novel series of 3,4,5-trisubstituted phenoxyacetic acid analogs. *Aust. J. Basic Appl. Sci.* 2007a;1(4):525-32.
- Dahiya R, Kaur R. Synthesis and anthelmintic potential of a novel series of 2-mercaptobenzimidazolepeptides. *Biosci. Biotech. Res. Asia* 2007b;4(2):561-6.
- Dahiya R, Kaur R. Synthesis of some 1,2,5-trisubstituted benzimidazole analogs as possible anthelmintic and antimicrobial agents. *Int. J. Biol. Chem. Sci.* 2008;2(1):1-13.
- Dahiya R, Kumar A. Synthesis, spectral and anthelmintic studies on some novel imidazole derivatives. *E-J. Chem.* 2008;5(S2):1133-43. [DOI: 10.1155/2008/161783]
- Dahiya R, Kumar A, Yadav R. Synthesis and biological activity of peptide derivatives of iodoquinazolinones/nitroimidazoles. *Molecules* 2008a;13(4):958-76. [DOI: 10.3390/molecules13040958]
- Dahiya R, Mourya R. Synthesis and antimicrobial screening of some novel halogenated phenoxyacetyl amino acid and peptide analogs. *Bull. Pharm. Res.* 2012;2(2):56-65.
- Dahiya R, Mourya R, Agrawal SC. Synthesis and antimicrobial screening of peptidyl derivatives of bromocoumarins/methyl imidazoles. *Afr. J. Pharm. Pharmacol.* 2010;4(5):214-25.
- Dahiya R, Pathak D. Synthesis and antimicrobial activity of novel series of 2-aryl-6-nitrobenzoyl amino acids and peptides. *Orient. J. Chem.* 2006a;22(1):123-8.
- Dahiya R, Pathak D. Synthesis and biological screening of a novel series of 6-nitro-2-(4-toluoyl)benzoyl amino acids and peptides. *J. Nepal Pharm. Assoc.* 2006b;XXIV(1):17-26.
- Dahiya R, Pathak D. Synthesis of novel heterocyclic analogs of 5-(4-methylcarboxamidophenyl)-2-furoic acid as potent antimicrobial agents. *Indian J. Heterocycl. Chem.* 2006c;16(1):53-6.
- Dahiya R, Pathak D. Cyclic peptides: New hope for antifungal therapy. *Egypt. Pharm. J. (NRC)* 2006d;5(2): 189-99.
- Dahiya R, Pathak D. Synthetic studies on novel benzimidazolepeptides with antimicrobial, cytotoxic and anthelmintic potential. *Eur. J. Med. Chem.* 2007;42(6):772-98. [DOI: 10.1016/j.ejmech.2006.11.015]
- Dahiya R, Pathak D, Bhatt S. Synthesis and biological evaluation of a novel series of 2-(4-chloro-3-methylphenoxy)acetyl amino acids and peptides. *J. Saudi Chem. Soc.* 2006a;10(1):165-76.
- Dahiya R, Pathak D, Bhatt S. Synthesis and biological evaluation of a novel series of 2-(2'-isopropyl-5'-methylphenoxy)acetyl amino acids and dipeptides. *Bull. Chem. Soc. Ethiop.* 2006b;20(2):235-45.
- Dahiya R, Pathak D, Kaur R. Synthesis, characterization and biological screening of some new aryloxyacetic acid analogs. *J. Indian Chem. Soc.* 2008b;85(7):754-8.
- de Moura RM, Pereira PS, Januário AH, França Sde C, Dias DA. Antimicrobial screening and quantitative determination of benzoic acid derivative of *Gomphrena celosioides* by TLC-densitometry. *Chem. Pharm. Bull. (Tokyo)* 2004;52(11):1342-4.
- Gao X, Cai X, Yan K, Song B, Gao L, Chen Z. Synthesis and antiviral bioactivities of 2-aryl- or 2-methyl-3-(substituted-benzalamino)-4-quinazolinone derivatives. *Molecules* 2007;12(12):2621-42. [DOI: 10.3390/12122621]
- Himaja M, Rajiv, Ramana MV. Synthesis of 6-nitrobenzimidazol-1-acetyl amino acids and peptides as potent anthelmintic agents. *Indian J. Heterocycl. Chem.* 2002; 12(2):121-4.

- Himaja M, Rajiv, Ramana MV, Poojary B, Satyanarayana D, Subrahmanyam EV, Bhat KI. Synthesis and biological activity of a novel series of 4-[2'-(6'-nitro)-benzimidazolyl]benzoyl amino acids and peptides. *Boll. Chim. Farm.* 2003;142(10):450-3.
- Khan ZK. In vitro and vivo screening techniques for bioactivity screening and evaluation, In: Proceeding Int. workshop UNIDO-CDRI, 1997; 210-1.
- Poojary B, Belagali SL, Kumar KH, Holla BS. Synthetic and antibacterial studies on some new furanopeptides. *Il Farmaco* 2003;58:569-72. [DOI: 10.1016/S0014-827X(03)00092-2]
- Poojary B, Belagali SL, Kumar KH, Holla BS, Bhagwat A. Synthesis and biological evaluation of amino acid derivatives of benzothiophenes. *Indian J. Heterocycl. Chem.* 2001;10:249-52.
- Sharma PC, Kaur G, Pahwa R, Sharma A, Rajak H. Quinazolinone analogs as potential therapeutic agents. *Curr. Med. Chem.* 2011;18(31):4786-812.
- Trusheva B, Todorov I, Ninova M, Najdenski H, Daneshmand A, Bankova V. Antibacterial mono- and sesquiterpene esters of benzoic acids from Iranian propolis. *Chem. Cent. J.* 2010;4(8):1-4. [DOI: 10.1186/1752-153X-4-8]
- Wu H, Xie X, Liu G. Parallel solution phase synthesis of 3,6,7-4(3H)-quinazolinones and evaluation of their antitumor activities against human cancer. *J. Comb. Chem.* 2010;12(3):346-55. [DOI: 10.1021/cc900173s]
- Wu J, Zhao M, Peng S. Quinazolin-4(3H)-ones capable of upregulating the expression of endogenous apolipoprotein A-1. *Expert Opin. Ther. Pat.* 2011;21(3):431-5. [DOI: 10.1517/13543776.2011.548322]
