

Maheshwari RK, Gupta S, Gharia A, Garg SK, Shilpkar R. Simple ecofriendly spectrophotometric estimation of tinidazole tablets by application of mixed-solvency techniques. *Bull. Pharm. Res.* 2011;1(1):22-5.

Cited by:

1. Jain R, Jain N, Jain DK, Jain SK. Eco-friendly quantitative estimation of lercanidipine hydrochloride: A novel approach using hydrotropic solubilization technique. *Bull. Pharm. Res.* 2013;3(1):23-8.
<http://www.appconnect.in/wp-content/uploads/2011/03/ReprintBPR067.pdf>
2. Pandey S, Pandey P, Dubey S, Chaturvedi U, Rai AK. Facile derivative UV spectroscopy method: simultaneous estimation of tinidazole and fluconazole in combined tablet dosage form. *Thai J. Pharm. Sci.* 2012;36(2):55-62.
<http://161.200.184.9/tjps/ContentVol36No2/V36-2%20Art2%20pp55-62.pdf>
3. Maheshwari RK, Shilpkar R. Formulation development and evaluation of injection of poorly soluble drug using mixed solvency concept. *Int. J. Pharm. Bio Sci.* 2012;3(1):179-89.
<http://ijpbs.net/vol-3/issue-1/bio/P%20-%2021.pdf>
4. Mehrotra A, Malviya G, Maheshwari RK. Application of mixed hydrotrophy in spectrophotometric analysis of frusemide in different formulations. *Bull. Pharm. Res.* 2011;1(2):15-21.
<http://www.appconnect.in/app/journalUploads/FirstPagePreviewBPR-2-4.pdf>

