

Pabreja K, Dua K. Comparative evaluation of *in situ* intestinal absorption of aceclofenac from solid dispersions,  $\beta$ -cyclodextrin complexes and co-precipitates in rats. *Bull. Pharm. Res.* 2011;1(1):26-30.

### References (15):

1. Anastasiadou C, Henry S, Legendre B, Souleau C, Duchene D. Solid dispersions: comparison of prepared melts and coprecipitates of diazepam and polyoxyethylene glycol 4000. *Drug Dev. Ind. Pharm.* 1983;9(1-2):103-15.  
<http://informahealthcare.com/toc/ddi/9/1-2>
2. Moffat AC, Osselton MD, Widdop B. Clarke's Analysis of Drugs and Poisons, Pharmaceutical Press, 1999;3:17-9.
3. Damian F, Blaton N, Kinget R, Van den Mooter G. Physical stability of solid dispersions of the antiviral agent UC-781 with PEG 6000, Gelucire® 44/14 and PVP K30. *Int. J. Pharm.* 2002;244(1-2):87-98.  
<http://cat.inist.fr/?aModele=afficheN&cpsid=13887847>
4. Dua K, Pabreja K, Ramana MV. Preparation, characterization and *in vitro* evaluation of aceclofenac solid dispersions. *ARS Pharmaceutica* 2010;51(1):57-76.  
<http://farmacia.ugr.es/ars/pdf/576.pdf>
5. Dua K, Ramana MV, Sara UVS, Himaja M, Agrawal A, Garg V, Pabreja K. Investigation of enhancement of solubility of norfloxacin  $\beta$ -cyclodextrin in presence of acidic solubilizing additives. *Curr. Drug Deliv.* 2007;4(1):21-5.  
<http://www.benthamscience.com/cdd/sample/cdd4-1/003AP.pdf>
6. Dua K, Sara UVS, Ramana MV. Studies on release of aceclofenac from cyclodextrin systems. *Int. J. Pharm. Excip.* 2007b;4:101-5.
7. Jun SW, Kim M-S, Kim J-S, Park HJ, Lee S, Woo J-S, Hwang S-J. Preparation and characterization of simvastatin/hydroxypropyl- $\beta$ -cyclodextrin inclusion using supercritical antisolvent (SAS) process. *Eur. J. Pharm. Biopharm.* 2007;66(3):413-21.  
<http://cat.inist.fr/?aModele=afficheN&cpsid=18817185>
8. Kumar N, Jain AK, Singh C, Kumar R. Development, characterization and solubility study of solid dispersion of terbinafine hydrochloride by solvent evaporation method. *Asian J. Pharm.* 2008;2(3):154-58.  
<http://www.asiapharmaceutics.info/article.asp?issn=0973-8398;year=2008;volume=2;issue=3;spage=154;epage=158;aulast=Kumar>
9. Martin A, Doluisio JT. Industrial bioavailability and pharmacokinetics: Guidelines, regulations, and controls. College of Pharmacy, Drug Dynamics Institute, University of Texas, Austin, TX 78712. 1977;539.  
<http://onlinelibrary.wiley.com/doi/10.1002/jps.2600671058/abstract>

10. Martin-Algarra RV, Pascual-Costa RM, Merino M, Casabo VG. Effects of polysorbate 80 on amiodarone intestinal absorption in the rat. *Int. J. Pharm.* 1995;122(1-2):1-8.  
<http://www.ingentaconnect.com/content/els/03785173/1995/00000122/00000001/art00025>
11. Patel HM, Suhagia BN, Shah SA, Rathod IS, Parmar VK. Preparation and characterization of etoricoxib- $\beta$ -cyclodextrin complexes prepared by the kneading method. *Acta Pharm.* 2007;57(3):351-9.  
<http://public.carnet.hr/acphee/35107.pdf>
12. Patil MP, Gaikwad NJ. Preparation and characterization of gliclazide-polyethylene glycol 4000 solid dispersions. *Acta. Pharm* 2009;59(1):57-65.  
<http://public.carnet.hr/acphee/5709.pdf>
13. Rawat S, Jain SK. Enhancement of intestinal absorption of few cox-2 inhibitors through interaction with  $\beta$ -cyclodextrin. *Indian J. Pharm. Sci.* 2007;69(4):529-34.  
[http://www.ijpsonline.com/temp/IndianJPharmSci694529-6333972\\_173539.pdf](http://www.ijpsonline.com/temp/IndianJPharmSci694529-6333972_173539.pdf)
14. Shah JC, Chen JR, Chow D. Metastable polymorph of etoposide with higher dissolution rate. *Drug Dev. Ind. Pharm.* 1999;25(1):63-7.  
<http://informahealthcare.com/doi/abs/10.1081/DDC-100102142?journalCode=ddi>
15. Zakeri-Milani P, Valizadeh H, Tajerzadeh H, Azarmi Y, Islambolchilar Z, Barzegar S, Barzegar-Jalali M. Predicting human intestinal permeability using single-pass intestinal perfusion in rat. *J. Pharm. Pharm. Sci.* 2007;10(3):368-379.  
[http://www.ualberta.ca/~csps/JPPS10\\_3/MS\\_1461/MS\\_1461.html](http://www.ualberta.ca/~csps/JPPS10_3/MS_1461/MS_1461.html)

