

Kumar M, Dureja H. Development and characterization of factorially designed 5-fluorouracil microspheres. *Bull. Pharm. Res.* 2011;1(1):54-61.

References (13):

1. Arshady R. Albumin microspheres and microcapsules: methodology of manufacturing techniques. *J. Control. Rel.* 1990;14(2):111-31.
<http://cat.inist.fr/?aModele=afficheN&cpsidt=19335069>
2. Choi HS, Seo SA, Khang G, Rhee JM, Lee HB. Preparation and characterization of fentanyl-loaded PLGA microspheres: *in vitro* release profiles. *Int. J. Pharm.* 2002;234(1-2):195-203.
<http://www.ncbi.nlm.nih.gov/pubmed/11839450>
3. Davis SS, Illum L. Polymeric microspheres as drug carriers. *Biomaterials* 1988;9(1):111-5.
<http://www.ncbi.nlm.nih.gov/pubmed/3280037>
4. Duarte ARC, Roy C, Vega-Gonzalez A, Duarte CMM, Subra-Paternault P. Preparation of acetazolamide composite microparticles by supercritical anti-solvent techniques. *Int. J. Pharm.* 2007;332(1-2):132-9.
<http://3bs.expertissues.org/download.php?file=321&type=4&doc=pubs>
5. Esposito E, Cortesi R, Nastruzzi C. Gelatin microspheres: influence of preparation parameters and thermal treatment on chemico-physical and biopharmaceutical properties. *Biomaterials* 1996;17(20):2009-20.
<http://www.ncbi.nlm.nih.gov/pubmed/8894096>
6. Esposito E, Menegatti E, Cortesi R. Hyaluronan-based microspheres as tools for drug delivery: a comparative study. *Int. J. Pharm.* 2005;288(1):35-49.
<http://www.ncbi.nlm.nih.gov/pubmed/15607256>
7. Kim BK, Hwang SJ, Park JB, Park HJ. Preparation and characterization of drug-loaded polymethacrylate microspheres by an emulsion solvent evaporation method. *J. Microencapsul.* 2002;19(6):811-22.
<http://www.ncbi.nlm.nih.gov/pubmed/12569029>
8. Lamprecht A, Ubrich N, Perez MH, Lehr C-M, Hoffman M, Maincent P. Influence of process parameters on nanoparticle preparation performed by a double emulsion pressure homogenization technique. *Int. J. Pharm.* 2000;196(2):177-82.
<http://www.ncbi.nlm.nih.gov/pubmed/10699713>
9. Lorenzo-Lamosa ML, Remunan-Lopez C, Vila-Jato JL, Alonso MJ. Design of microencapsulated chitosan microspheres for colonic drug delivery. *J. Control. Rel.* 1998;52(1-2):109-18.
<http://www.ncbi.nlm.nih.gov/pubmed/9685941>

10. Murakami H, Kobayashi M, Takeuchi H, Kawashima Y. Further application of a modified spontaneous emulsification solvent diffusion method to various types of PLGA and PLA polymers for preparation of nanoparticles. *Powder Tech.* 2000;107(1-2):137-43.
<http://cat.inist.fr/?aModele=afficheN&cpsidt=1260785>
11. Ritschel WA. Biopharmaceutic and pharmacokinetic aspects in the design of controlled release peroral drug delivery systems. *Drug Dev. Ind. Pharm.* 1989;15(6-7):1073-1103.
<http://informahealthcare.com/doi/abs/10.3109/03639048909043666?journalCode=ddi>
12. Sahoo SK, Mallick AA, Barik BB, Senapati, PC. Formulation and *in vitro* evaluation of eudragit® microspheres of stavudine. *Trop. J. Pharm. Res.* 2005;4(1):369-75.
http://www.tjpr.org/vol4_no1/41sahoo.pdf
13. Yang Y-Y, Chung T-S, Ng NP. Morphology, drug distribution, and *in vitro* release profiles of biodegradable polymeric microspheres containing protein fabricated by double-emulsion solvent extraction/evaporation method. *Biomaterials* 2001;22(3):231-41.
<http://www.ncbi.nlm.nih.gov/pubmed/11197498>

