

Sachan NK, Pushkar S. Solid dispersions: An industrially feasible alternative approach to formulate brick dust molecules. *Bull. Pharm. Res.* 2011;1(1):75-87.

References (59):

1. Aleem MA. Solid dispersion - An approach to enhance the dissolution rate of aceclofenac. Rajiv Gandhi University of Health Sciences, Bangalore, Karnatka (India). M.Pharm thesis, 2006;1-24.
2. Aungst BJ, Nguyen NH, Rogers NJ, Rowe SM, Hussain MA, White SJ, Shum L. Amphiphilic vehicles improve the oral bioavailability of a poorly soluble HIV protease inhibitor at high doses. *Int. J. Pharm.* 1997;156(1):79-88.
<http://www.sciencedirect.com/science/article/pii/S0378517397001890>
3. Betageri GV, Makarla KR. Enhancement of dissolution of glyburide by solid dispersion and lyophilization techniques. *Int. J. Pharm.* 1995;126(1-2):155-60.
<http://www.sciencedirect.com/science/article/pii/0378517395041141>
4. Beten DB, Amighi K, Moes AJ. Preparation of controlled-release coevaporates of dipyridamole by loading neutral pellets in a fluidized-bed coating system. *Pharm. Res.* 1995;12(9):1269-72.
<http://164.15.112.38/ULB-GALENIQUE/pdf/Publications/Beten%20et%20al%20-%20PharmRes%201995.pdf>
5. Breitenbach J. Melt extrusion: from process to drug delivery technology. *Eur. J. Pharm. Biopharm.* 2002;54(2):107-17.
<http://www.ncbi.nlm.nih.gov/pubmed/12191680>
6. Buckton G, Darcy P. The use of gravimetric studies to assess the degree of crystallinity of predominantly crystalline powders. *Int. J. Pharm.* 1995;123(2):265-71.
<http://cat.inist.fr/?aModele=afficheN&cpsidt=3615785>
7. Chatham SM. The use of bases in SSM formulations. *STP Pharm.* 1987;3:575-82.
8. Chiou WL, Riegelman S. Pharmaceutical applications of solid dispersion systems. *J. Pharm. Sci.* 1971;60(9):1281-1302.
<http://onlinelibrary.wiley.com/doi/10.1002/jps.2600600902/abstract>
9. Chiou WL, Smith LD. Solid dispersion approach to the formulation of organic liquid drugs using polyethylene glycol 6000 as a carrier. *J. Pharm. Sci.* 1971;60(1):125-7.
<http://onlinelibrary.wiley.com/doi/10.1002/jps.2600600127/abstract>
10. Cilurzo F, Minghetti P, Casiraghi A, Montanari L. Characterization of nifedipine solid dispersions. *Int. J. Pharm.* 2002;242(1-2):313-7.
<http://www.sciencedirect.com/science/article/pii/S0378517302001734>
11. Corrigan OI. Mechanisms of dissolution of fast release solid dispersions. *Drug Dev. Ind. Pharm.* 1985;11(2-3):697-724.
<http://informahealthcare.com/doi/abs/10.3109/03639048509056896>

12. Corrigan OI. Retardation of polymeric carrier dissolution by dispersed drugs: Factors influencing the dissolution of solid dispersions containing polyethylene glycols. *Drug Dev. Ind. Pharm.* 1986;12(11-13):1777-93.
<http://informahealthcare.com/doi/abs/10.3109/03639048609042609?journalCode=ddi>
13. Daabis NA, Abd-Elfattah S, el-Banna HM. Physicochemical study of drug binary systems. 2. Khellin-urea system, *Pharmazie* 1974;29(6):400-4.
<http://www.ncbi.nlm.nih.gov/pubmed/4853639>
14. De Meuter P, Rahier H, Van Mele B. The use of modulated temperature differential scanning calorimetry for the characterisation of food systems. *Int. J. Pharm.* 1999;192(1):77-84.
<http://www.sciencedirect.com/science/article/pii/S0378517399002744>
15. Dennis AB, Farr SJ, Kellaway IW, Taylor G, Davidson R. In vivo evaluation of rapid release and sustained release Gelucire capsule formulations. *Int J Pharm.* 1990;65(1-2):85-100.
<http://www.sciencedirect.com/science/article/pii/037851739090013T>
16. Dhirendra K, Lewis S, Udupa N, Atin K. Solid dispersions: a review. *Pak. J. Pharm. Sci.* 2009;22(2):234-46.
<http://www.pjps.pk/CD-PJPS-22-2-09/Paper-20.pdf>
17. Dordunoo SK, Ford JL, Rubinstein MH. Preformulation studies on solid dispersions containing triamterene or temazepam in polyethylene glycols or Gelucire 44/14 for liquid filling of hard gelatin capsules. *Drug Dev. Ind. Pharm.* 1991;17(12):1685-1713.
<http://informahealthcare.com/doi/abs/10.3109/03639049109057315>
18. Dua K, Pabreja K, Sharma VK, Sara UVS, Ramana MV. Solid dispersion technology. *Pharmabiz* 2009; September 10, Thursday.
<http://saffron.pharmabiz.com/article/detnews.asp?Arch=&articleid=51592§ionid=46>
19. Dubois J-L, Ford JL. Similarities in the release rates of different drugs from polyethylene glycol 6000 solid dispersions. *J. Pharm. Pharm.* 1985;37(7):494-95.
<http://onlinelibrary.wiley.com/doi/10.1111/j.2042-7158.1985.tb03048.x/abstract>
20. El-Badry M, Fathy M. Enhancement of the dissolution and permeation rates of meloxicam by formation of its freeze-dried solid dispersions in polyvinylpyrrolidone K-30. *Drug Dev. Ind. Pharm.* 2006;32(2):141-50.
<http://www.ncbi.nlm.nih.gov/pubmed/16537195>
21. Fathy M, Sheha M. *In vitro* and *in vivo* evaluation of an amylobarbitone/hydroxypropyl-beta-cyclodextrin complex prepared by a freeze-drying method. *Pharmazie* 2000; 55(7):513-7.
<http://www.ncbi.nlm.nih.gov/pubmed/10944779>
22. Ford JL, Rubinstein MH. Phase equilibria and dissolution rates of indomethacin-polyethylene glycol 6000 solid dispersions. *Pharm. Acta Helv.* 1978;53(11):327-32.
<http://www.ncbi.nlm.nih.gov/pubmed/746042>
23. Ghaderi R. A supercritical fluids extraction process for the production of drug loaded biodegradable microparticles. *Acta Universitatis Upsaliensis*, Ph.D Thesis, Uppsala University, Sweden, 2000;1-46.
<http://www.openthesis.org/documents/supercritical-fluids-extraction-process-production-412309.html>

24. Gines JM, Veiga MD, Arias MJ, Rabasco AM. Elaboration and thermal study of interactions between cinnarizine and gelucire® 53/10 physical mixtures and solid dispersions. *Int. J. Pharm.* 1995;126(1-2):287-91.
<http://www.sciencedirect.com/science/article/pii/0378517395041400>
25. Goldberg AH, Gibaldi M, Kanig JL. Increasing dissolution rates and gastrointestinal absorption of drugs *via* solid solutions and eutectic mixtures II: Experimental evaluation of a eutectic mixture: Urea-acetaminophen system. *J. Pharm. Sci.* 1966;55(5):482-7.
<http://onlinelibrary.wiley.com/doi/10.1002/jps.2600550507/abstract>
26. Higuchi WI. Diffusional models useful in biopharmaceutics. Drug release rate processes. *J. Pharm. Sci.* 1967;56(3):315-24.
<http://onlinelibrary.wiley.com/doi/10.1002/jps.2600560302/abstract>
27. Hite M, Turner S, Federici C. Part 1: Oral delivery of poorly soluble drugs. *Pharmaceutical Manufacturing and Packing Sourcer* Summer 2003;03:1-3.
http://s15611.gridserver.com/lit/PMPS_2003_1.pdf
28. Ho H-O, Su H-L, Tsai T, Sheu M-T. The preparation and characterization of solid dispersions on pellets using a fluidized-bed system. *Int. J. Pharm.* 1996;139(1-2):223-9.
<http://www.sciencedirect.com/science/article/pii/0378517396045942>
29. Hulsmann S, Backensfeld T, Keitel S, Bodmeier R. Melt extrusion-an alternative method for enhancing the dissolution rate of 17 β -estradiol hemihydrate. *Eur. J. Pharm. Biopharm.* 2000;49(3):237-42.
<http://www.sciencedirect.com/science/article/pii/S0939641100000771>
30. Jovanovic N, Bouchard A, Hofland GW, Witkamp G-J, Crommelin DJA, Jiskoot W. Stabilization of proteins in dry powder formulations using supercritical fluid technology. *Pharm. Res.* 2004;21(11):1955-69.
<http://www.springerlink.com/content/q141467442837010/>
31. Kang BK, Lee JS, Chon SK, Jeong SY, Yuk SH, Khang G, Lee HB, Cho SH. Development of self-microemulsifying drug delivery systems (SMEDDS) for oral bioavailability enhancement of simvastatin in beagle dogs. *Int. J. Pharm.* 2004;274(1-2):65-73.
<http://www.sciencedirect.com/science/article/pii/S0378517304000043>
32. Karavas E, Ktistis G, Xenakis A, Georgarakis E. Effect of hydrogen bonding interactions on the release mechanism of felodipine from nanodispersions with polyvinylpyrrolidone. *Eur. J. Pharm. Biopharm.* 2006;63(2):103-14.
<http://www.sciencedirect.com/science/article/pii/S0939641106000476>
33. Kaushal AM, Gupta P, Bansal AK. Amorphous drug delivery systems: molecular aspects, design and performance. *Crit. Rev. Ther. Drug Carrier Syst.* 2004;21(3):133-93.
<http://www.begellhouse.com/journals/3667c4ae6e8fd136,16e2a9f911dc5f33,4c11e55e55217865.html>
34. Kennedy JP, Neibergall PJ. Development and optimization of a solid dispersion hot-melt fluid bed coating method. *Pharm. Dev. Technol.* 1996;1(1):51-62.
<http://www.ncbi.nlm.nih.gov/pubmed/9552331>
35. Leuner C, Dressman J. Improving drug solubility for oral delivery using solid dispersions. *Eur. J. Pharm. Biopharm.* 2000;50(1):47-60.
<http://www.sciencedirect.com/science/article/pii/S093964110000076X>

36. Lloyd GR, Craig DQM, Smith A. A calorimetric investigation into the interaction between paracetamol and polyethylene glycol 4000 in physical mixes and solid dispersions. *Eur. J. Pharm. Biopharm.* 1999;48(1):59-65.
<http://www.sciencedirect.com/science/article/pii/S0939641199000223>
37. Muhrer G, Meier U, Fusaro F, Albano S, Mazzotti M. Use of compressed gas precipitation to enhance the dissolution behavior of a poorly water-soluble drug: Generation of drug microparticles and drug-polymer solid dispersions. *Int. J. Pharm.* 2006;308(1-2):69-83.
<http://www.sciencedirect.com/science/article/pii/S037851730500709X>
38. Perissutti B, Newton JM, Podcizek F, Rubessa F. Preparation of extruded carbamazepine and PEG 4000 as a potential rapid release dosage form. *Eur. J. Pharm. Biopharm.* 2002;53(1):125-32.
<http://www.sciencedirect.com/science/article/pii/S0939641101002090>
39. Phillips EM, Stella VJ. Rapid expansion from supercritical solutions: Applications to pharmaceutical processes. *Int. J. Pharm.* 1993;94(1-3):1-10.
<http://www.sciencedirect.com/science/article/pii/037851739390002W>
40. Pokharkar VB, Mandpe LP, Padamwar MN, Ambike A, Mahadik KR, Paradkar A. Development, characterization and stabilization of amorphous form of low Tg drug. *Powder Technol.* 2006;167(1):20-5.
<http://www.sciencedirect.com/science/article/pii/S0032591006001732>
41. Reneker DH, Chun I. Nanometer diameter fibers of polymers produced by electro-spinning. *Nanotechnology* 1996;7(3):216-23.
<http://iopscience.iop.org/0957-4484/7/3/009;jsessionid=ED47387D4A664D562173546627174A47.c3>
42. Sebhatu T, Angberg M, Ahlneck C. Assessment of the degree of disorder in crystalline solids by isothermal microcalorimetry. *Int. J. Pharm.* 1994;104(2):135-44.
<http://www.sciencedirect.com/science/article/pii/0378517394901880>
43. Sekiguchi K, Obi N. Studies on absorption of eutectic mixture. I. A comparison of the behavior of eutectic mixture of sulphathiazole and that of ordinary sulphathiazole in man. *Chem. Pharm. Bull.* 1961;9(11):866-72.
<http://www.journalarchive.jst.go.jp/jnlpdf.php?cdjournal=cpb1958&cdvol=9&noissue=11&startpage=866&lang=en&from=jnlto>
44. Sekiguchi K, Obi N, Ueda Y. Studies on absorption of eutectic mixture. II. Absorption of fused conglomerates of chloramphenicol and urea in rabbits. *Chem. Pharm. Bull.* 1964;12(2):134-44.
<http://www.journalarchive.jst.go.jp/jnlpdf.php?cdjournal=cpb1958&cdvol=12&noissue=2&startpage=134&lang=en&from=jnlto>
45. Serajuddin ATM. Bioavailability enhancement of poorly water-soluble drugs by solid dispersion in surface-active vehicles: A recent breakthrough. *La Gazette de l'ApGI (France)* 1997a;14:144-5.
46. Serajuddin ATM. Solid dispersion of poorly water-soluble drugs: Early promises, subsequent problems, and recent breakthroughs. *J. Pharm. Sci.* 1999;88(10):1058-66.
<http://onlinelibrary.wiley.com/doi/10.1021/js980403l/abstract>

47. Serajuddin ATM, Li P, Haefele TF. Development of lipid-based drug delivery systems for poorly water-soluble drugs as viable oral dosage forms - present status and future prospects. *Am. Pharm. Rev.* 2008;11(2):34-42.
<http://americanpharmaceuticalreview.com/ViewArticle.aspx?ContentID=3378>
48. Serajuddin ATM. Bioavailability enhancement of poorly water soluble drugs by solid dispersions in surface active and self-emulsifying vehicles. *Bull. Techniq. Gattefosse* 1997b;90:43-50.
49. Serajuddin ATM, Sheen P-C, Mufson D, Bernstein DF, Augustine MA. Effect of vehicle amphiphilicity on the dissolution and bioavailability of a poorly water-soluble drug from solid dispersions.
<http://onlinelibrary.wiley.com/doi/10.1002/jps.2600770512/abstract>
50. Serajuddin ATM, Sheen P-C, Mufson D, Bernstein DF, Augustine MA. Physicochemical basis of increased bioavailability of a poorly water-soluble drug following oral administration as organic solutions. *J. Pharm. Sci.* 1988b;77(4):325-9.
<http://onlinelibrary.wiley.com/doi/10.1002/jps.2600770409/abstract>
51. Squillante E, Neduvelil S. Preparation of sustained-release chlorpheniramine maleate powders using supercritical fluid processing technology. *AAPS Annual Meeting*, November 14-18, 1999; Poster 3617.
http://www.aapsi.org/abstracts/AM_1999/957.htm
52. Tachibana T, Nakamura A. A method for preparing an aqueous colloidal dispersion of organic materials by using water-soluble polymers: dispersion of β -carotene by polyvinylpyrrolidone. *Colloid. Polym. Sci.* 1965;203(2):130-3.
<http://www.springerlink.com/content/jx6v4n1k33783831/>
53. Taylor LS, Zografi G. Spectroscopic characterization of interactions between PVP and indomethacin in amorphous molecular dispersions. *Pharm. Res.* 1997;14(12):1691-8.
<http://www.springerlink.com/content/p2v6906166147229/>
54. Thakkar AL, Hirsch CA, Page JG. Solid dispersion approach for overcoming bioavailability problems due to polymorphism of nabilone, a cannabinoid derivative. *J. Pharm. Pharmacol.* 1977;29(1):783-4.
<http://onlinelibrary.wiley.com/doi/10.1111/j.2042-7158.1977.tb11468.x/abstract>
55. Tsinontides SC, Rajniak P, Pham D, Hunke WA, Placek J, Reynolds SD. Freeze drying-principles and practice for successful scale-up to manufacturing. *Int. J. Pharm.* 2004;280(1-2):1-16.
<http://www.sciencedirect.com/science/article/pii/S0378517304002479>
56. van Drooge D-J, Hinrichs WL, Dickhoff BH, Elli MN, Visser MR, Zijlstra GS, Frijlink HW. Spray freeze drying to produce a stable Delta(9)-tetrahydrocannabinol containing inulin-based solid dispersion powder suitable for inhalation. *Eur. J. Pharm. Sci.* 2005;26(2):231-40.
<http://www.sciencedirect.com/science/article/pii/S0928098705001995>
57. Vasanthavada M, Tong W-Q, Joshi Y, Kislalioglu MS. Phase behavior of amorphous molecular dispersions I: Determination of the degree and mechanism of solid solubility. *Pharm. Res.* 2004;21(9):1598-1606.
<http://www.springerlink.com/content/x577387xu0107126/>

58. Verreck G, Chun I, Peeters J, Rosenblatt J, Brewster ME. Preparation and characterization of nanofibers containing amorphous drug dispersions generated by electrostatic spinning. *Pharm. Res.* 2003;20(5):810-7.
<http://www.springerlink.com/content/t831m0727hw83646/>
59. Willard HH, Merritt Jr LL, Dean JA. Instrumental methods of Analysis, 4th edition, Van Nostrand Reinhold Company: New York, 1965; 198-238.

