

Shrivastava J, Shah K, Mahadik M, Dhaneshwar SR. Application of HPTLC in the simultaneous estimation of thiocolchicoside and diclofenac in bulk drug and pharmaceutical dosage form. *Bull. Pharm. Res.* 2011;1(3):34-7.

References (13):

1. Balduini W, Cimino M, Depoortere H, Cattabeni F. Characterization of [³H]thiocolchicoside binding sites in rat spinal cord and cerebral cortex. *Eur. J. Pharmacol.* 1999; 376(1-2):149-157.
<http://www.sciencedirect.com/science/article/pii/S0014299999003714>
2. British Pharmacopoeia, Volume II, British Pharmacopoeial Commission, Her Majesty Stationery Office: London, 2003; 582-3.
3. Cimino M, Marini P, Cattabeni F. Interaction of thiocolchicoside with [³H]strychnine binding sites in rat spinal cord and brainstem. *Eur. J. Pharmacol.* 1996; 318(1):201-4.
<http://www.sciencedirect.com/science/article/pii/S0014299996008849>
4. El-Ragehy NA, Ellaithy MM, El-Ghobashy MA. Determination of thiocolchicoside in its binary mixtures (thiocolchicoside-glafenine and thiocolchicoside-floctafenine) by TLC-densitometry. *Farmaco* 2003;58(6):463-8.
<http://www.ncbi.nlm.nih.gov/pubmed/12767387>
5. Ferrari MP, Gatti G, Fattore C, Fedele G, Novellini R. Comparative bioavailability and tolerability study of two intramuscular formulations of thiocolchicoside in healthy volunteers. *Eur. J. Drug Metab. Pharmacokinet.* 2001;26(4):257-62.
<http://www.ncbi.nlm.nih.gov/pubmed/11808868>
6. ICH, Q2 (R1). Validation of Analytical Procedures: Text and Methodology. International Conference on Harmonisation, Geneva, 2005.
7. Patil MG, Banerjee SK, Bonde CG, Chhabra GS. UV spectrophotometric method development for the determination of desvenlafaxine succinate in tablet formulation. *Bull. Pharm. Res.* 2011;1(1):40-3.
<http://www.appconnect.in/wp-content/uploads/2011/05/FirstPagePreviewBPR007.pdf>
8. Prasanthi V, Mary K, Narasimha Raju CH, Basaveswara Rao MV. Development and validation of new RP-HPLC method for determination of acetyl sulfisoxazole in bulk and pharmaceutical dosage forms. *Bull. Pharm. Res.* 2011;1(1):47-53.
<http://www.appconnect.in/wp-content/uploads/2011/05/FirstPagePreviewBPR009.pdf>
9. Rosso A, Zuccaro S. Determination of alkaloids from the colchicine family by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A* 1998;825(1):96-101.
<http://www.sciencedirect.com/science/article/pii/S0021967398005950>

10. Shah J, Banerjee SK, Chhabra GS. UV spectrophotometric method development and validation for entacapone in bulk and formulation. *Bull. Pharm. Res.* 2011;1(2):7-9.
<http://www.appconnect.in/app/journalUploads/FirstPagePreviewBPR-2-2.pdf>
11. Shukla R, Shivkumar R, Shivan KN. Development of a UV-spectrophotometric method for the simultaneous determination of tramadol hydrochloride and paracetamol in bulk and marketed product. *Bull. Pharm. Res.* 2011;1(1):62-6.
<http://www.appconnect.in/wp-content/uploads/2011/05/FirstPagePreviewBPR011.pdf>
12. Sutherland FCW, Smit MJ, Herbst L, Els J, Hundt HKL, Swart KJ, Hundt AF. Highly specific and sensitive liquid chromatography-tandem mass spectrometry method for the determination of 3-desmethylthiocolchicine in human plasma as analyte for the assessment of bioequivalence after oral administration of thiocolchicoside. *J. Chromatogr. A* 2002; 949(1-2):71-7.
<http://www.sciencedirect.com/science/article/pii/S0021967302000080>
13. Vargas F, Mendez H, Fuentes A, Sequera J, Fraile G, Velasquez M, Caceres G, Cuello K. Photosensitizing activity of thiocolchicoside: photochemical and in vitro photo-toxicity studies. *Pharmazie* 2001;56(1):83-8.
<http://www.ncbi.nlm.nih.gov/pubmed/11210677>

